## NATIONAL NUCLEAR DATA CENTER Bldg. 197D Brookhaven National Laboratory P. O. Box 5000 Upton, NY 11973-5000 U.S.A.

(Internet) "NNDC@BNL.GOV

Telephone: (516)344-2902 FAX: (516)344-2806

## Memo CP-C/349

DATE: August 11, 2004
TO: Distribution
FROM: V. McLane

**SUBJECT**: Particle specification and long reaction strings.

For DA/DA and DA/DA/DE data, I have come up with a fairly long reaction string. I can foresee in the future the strings getting even longer, especially in the compilation of heavy ion reactions and radioactive ion beam data.

<u>Problem 1</u>: When the particles considered are heavy ions, and neither is the residual nucleus, a particle code must be given in SF7. I propose that shortened versions of the nuclide code be used, for example, BE10 instead of 5-BE-10. This saves us 3 or 4 characters per code, and will put off the time when we have to deal with problem 2).

```
(3-LI-7(3-LI-7,2A)2-HE-6,PAR,DA/DA,BE10/A+HE6,REL)
```

<u>Problem 2</u>: With the compilation of heavy ion data, the reaction codes can easily exceed 55 characters. Again, this can be solved in the short time by allowing shortened versions of the nuclide code in SF2 and SF3. However, this may not be a long term solution as the quantities could still exceed 55 characters. A completely fictitious example with <u>75</u> characters is:

```
(29-CU-63(3-LI-7,2-HE-6+3-LI-6)27-CO-58,,DA/DA/DE,2-HE-6/3-LI-6/3-LI-6,REL)
```

this would be cut down to the following with  $\underline{66}$  characters using the solution given for problem 1).

(29-CU-63(3-LI-7, 2-HE-6+3-LI-6)27-CO-68,,DA/DA/DE,HE6/LI6/LI6,REL)

## Remark by O.Schwerer:

Possible solutions for "too long" reactions strings (More than one option may be implemented):

- Shortened nuclide code in SF7 and potentially also in SF2 and SF3 as proposed above
- Remember using RSD in SF7 when appropriate
- Perhaps introduce new code for compound nucleus, used for SF7
- Continuation records for REACTION codes