

Visit the [Isotope Explorer](#) home page!

14 reference(s) found :

Keynumber: 1995MO40

Reference: Aust.J.Phys. 48, 125 (1995)

Authors: A.J.Morton, D.G.Sargood

Title: Thermonuclear Reactions Rates for Reactions Leading to N = 28 Nuclei

Keyword abstract: NUCLEAR REACTIONS $^{44, 46}\text{K}$, $^{46, 47, 48}\text{Ca}$, $^{45, 47, 48, 49, 50}\text{Sc}$, $^{46, 47, 48, 49, 50}\text{Ti}$, $^{47, 48, 49, 50, 51}\text{V}$, $^{48, 49, 50, 51, 52}\text{Cr}$, $^{51, 52, 53}\text{Mn}$, $^{52, 53, 54}\text{Fe}$, $^{55}\text{Co}(n,\gamma)$, (n,p) , (n,α) , (p,γ) , (p,n) , (p,α) , (α,γ) , (α,n) , (α,p) , E not given; $^{56}\text{Ni}(n,\gamma)$, (n,p) , (n,α) , (α,γ) , (α,n) , (α,p) , E not given; ^{46}Ar , $^{45, 47}\text{K}$ (p,γ) , (p,n) , (p,α) , (α,γ) , (α,n) , (α,p) , E not given; calculated stellar reaction rates vs temperature. Statistical model calculations, optical-model potential.

Keynumber: 1989CO01

Reference: J.Phys.(London) G15, 321 (1989)

Authors: S.P.Collins, S.A.Eid, S.A.Hamada, W.D.Hamilton, F.Hoyler

Title: A Search for Mixed-Symmetry States in the Mass A \approx 50 Region

Keyword abstract: RADIOACTIVITY $^{56}\text{Mn}(\beta^-)$; measured $\gamma(\theta)$. ^{56}Fe levels deduced δ . Cryogenically oriented nuclei.

Keyword abstract: NUCLEAR REACTIONS ^{47}Ti , ^{53}Cr , $^{57}\text{Fe}(n,\gamma)$, E=thermal; measured $\gamma(\theta)$. ^{48}Ti , ^{54}Cr , ^{58}Fe levels deduced $\delta, \mu, B(\lambda)$. Enriched target, on-line directional correlations.

Keynumber: 1984RU06

Reference: Nucl.Phys. A419, 439 (1984)

Authors: J.F.A.G.Ruyl, J.B.M.De Haas, P.M.Endt, L.Zybert

Title: Investigation of the $^{47, 49}\text{Ti}(n,\gamma)^{48, 50}\text{Ti}$ Reactions

Keyword abstract: NUCLEAR REACTIONS $^{49, 47}\text{Ti}(n,\gamma)$, (polarized n, γ), E=thermal; measured $E\gamma, I\gamma$ ($E\gamma, \theta$), γ CP; deduced Q. $^{48, 50}\text{Ti}$ deduced levels, γ -branching, J, π . Enriched, polarized, unpolarized targets.

Keynumber: 1983SA30

Reference: Aust.J.Phys. 36, 583 (1983)

Authors: D.G.Sargood

Title: Effect of Excited States on Thermonuclear Reaction Rates

Keyword abstract: NUCLEAR REACTIONS, ICPND $^{20, 21, 22}\text{Ne}$, ^{23}Na , $^{24, 25, 26}\text{Mg}$, ^{27}Al , $^{28, 29, 30}\text{Si}$, ^{31}P , $^{32, 33, 34, 36}\text{S}$, $^{35, 37}\text{Cl}$, $^{36, 38, 40}\text{Ar}$, $^{39, 40, 41}\text{K}$, $^{40, 42, 43, 44, 46, 48}\text{Ca}$, ^{45}Sc , $^{46, 47, 48, 49, 50}\text{Ti}$, $^{50, 51}\text{V}$, $^{50, 52, 53, 54}\text{Cr}$, ^{55}Mn , $^{54, 56, 57, 58}\text{Fe}$, ^{59}Co , $^{58, 60, 61, 62, 64}\text{Ni}$, $^{63, 65}\text{Cu}$, $^{64, 66, 67}\text{Zn}(n,\gamma)$, (n,p) , (n,α) , (p,γ) , (p,n) , (p,α) , (α,γ) , (α,n) , (α,p) , $^{70}\text{Zn}(p,\gamma)$, (p,n) , (p,α) , (α,γ) , (α,n) , (α,p) , E=low; compiled target thermal distribution energy state to ground state thermonuclear reaction rate of reaction σ vs temperature. Statistical model.

Keynumber: 1980PIZN

Coden: CONF Kiev(Neutron Physics) Proc, Part3, P270, Pisanko

Keyword abstract: NUCLEAR REACTIONS $^{22, 23}\text{Na}$, $^{24, 25, 26}\text{Mg}$, ^{27}Al , $^{28, 29, 30}\text{Si}$, ^{31}P , $^{32, 33, 34}\text{S}$, $^{35, 36, 37}\text{Cl}$, $^{36, 38, 40}\text{Ar}$, $^{39, 40, 41}\text{K}$, $^{40, 42, 43, 44, 46, 48}\text{Ca}$, $^{45, 46}\text{Sc}$, $^{46, 47, 48, 49, 50}\text{Ti}$, $^{50, 51}\text{V}$, $^{50, 52, 53, 54}\text{Cr}$, $^{54, 56, 57, 58}\text{Fe}$, ^{59}Co , $^{58, 59, 60, 61, 62, 64}\text{Ni}$, $^{63, 65}\text{Cu}$, $^{64, 66, 67, 68, 70}\text{Zn}$, $^{69, 71}\text{Ga}(n,\gamma)$, (n,n) , (n,α) , E=thermal; evaluated σ , radiative capture

resonance integrals.

Keynumber: 1980IS02

Reference: Can.J.Phys. 58, 168 (1980)

Authors: M.A.Islam, T.J.Kennett, S.A.Kerr, W.V.Prestwich

Title: A Self-Consistent Set of Neutron Separation Energies

Keyword abstract: NUCLEAR REACTIONS ^1H , ^9Be , ^{14}N , 24 , ^{25}Mg , ^{27}Al , 28 , ^{29}Si , ^{32}S , ^{35}Cl , 40 , ^{44}Ca , 47 , 48 , ^{49}Ti , 50 , 52 , ^{53}Cr , ^{55}Mn , 54 , 56 , $^{57}\text{Fe}(n,\gamma)$, E=thermal; measured $E\gamma$, $I\gamma$. ^2H , ^{10}Be , 25 , ^{26}Mg , ^{28}Al , 29 , ^{30}Si , ^{33}S , ^{36}Cl , 41 , ^{45}Ca , 48 , 49 , ^{50}Ti , 51 , 53 , ^{54}Cr , ^{56}Mn , 55 , 57 , ^{58}Fe deduced Q,neutron binding energy.

Keynumber: 1979THZW

Reference: Proc.Specialsts Meeting on Neutron Data Structural Materials for Fast Reactors, December 5-8, 1977, Geel, Belgium, p.675 (1979)

Authors: B.Thom, D.B.Gayther, M.C.Moxon, B.W.Thomas

Title: Capture Cross-Section Measurements on the Separated Isotopes of Titanium

Keyword abstract: NUCLEAR REACTIONS 46 , 47 , 49 , $^{50}\text{Ti}(n,\gamma)$, E=low; measured capture σ . 47 , 48 , 50 , ^{51}Ti deduced resonance parameters.

Keynumber: 1977ALYR

Reference: AAEC/E-402 (1977)

Authors: B.J.Allen, J.W.Boldeman, A.R.de L.Musgrove, R.L.Macklin

Title: Resonance Neutron Capture in the Isotopes of Titanium

Keyword abstract: NUCLEAR REACTIONS 46 , 47 , 48 , 49 , $^{50}\text{Ti}(n,\gamma)$, E=2.75-300 keV; measured capture γ -yield. 47 , 48 , 49 , 50 , ^{51}Ti deduced resonance parameters.

Keynumber: 1971NEZZ

Coden: CONF Moscow(NuclSpectros,Structure) Abstr P38

Keyword abstract: NUCLEAR REACTIONS 46 , 47 , 48 , 49 , $^{50}\text{Ti}(n,\gamma)$, E not given; measured $E\gamma$, $I\gamma$. 47 , 48 , 49 , 50 , ^{51}Ti deduced transitions.

Keynumber: 1971ARZJ

Coden: CONF Legnaro($1f_{7/2}$ Nuclei),P251

Keyword abstract: NUCLEAR REACTIONS ^{36}Ar , ^{40}Ar , ^{40}K , 40 , 42 , 44 , 46 , ^{48}Ca , ^{47}Ti , ^{55}Mn , ^{57}Fe , $^{59}\text{Co}(n,\gamma)$, E=thermal; surveyed $E\gamma$, $I\gamma$, $\gamma\gamma$ -coin, $\gamma\gamma(\theta)$, γ -polarization data. ^{37}Ar , ^{41}Ar , ^{41}K , 41 , 43 , 45 , 47 , ^{49}Ca , ^{48}Ti , ^{56}Mn , ^{58}Fe , ^{60}Co deduced levels, J, π , γ -mixing.

Keynumber: 1969TE06

Reference: Phys.Rev. 187, 1403 (1969)

Authors: J.Tenenbaum, R.Moreh, Y.Wand, G.Ben-David

Title: Study of the Level Structure of ^{48}Ti Using the $^{47}\text{Ti}(n,\gamma)$ Reaction

Keyword abstract: NUCLEAR REACTIONS $^{47}\text{Ti}(n,\gamma)$, E=thermal; measured $E\gamma$, $I\gamma$, $\gamma\gamma(\theta)$; deduced Q. ^{48}Ti deduced levels, J, π , γ -mixing.

Keynumber: 1969FE08

Reference: Nucl.Phys. A139, 113 (1969)

Authors: P.Fettweis, M.Saidane

Title: The Level Scheme of ^{48}Ti and ^{49}Ti as Studied by the Neutron Capture γ -Ray Spectra

Keyword abstract: NUCLEAR REACTIONS $^{47, 48}\text{Ti}(n, \gamma)$, E= thermal; measured $E\gamma$, $I\gamma$. $^{48, 49}\text{Ti}$ deduced levels. Enriched ^{47}Ti target.

Keynumber: 1968BAZZ

Reference: Program and Theses, Proc.18th Ann.Conf.Nucl.Spectroscopy and Struct.Of At.Nuclei, Riga, p.32 (1968)

Authors: I.F.Barchuk, D.A.Bazavov, G.V.Belykh, V.I.Golyshkin, A.V.Murzin, A.F.Ogorodnik

Title: Spectra of γ -Rays Caused by Capture of Slow Neutrons by ^{25}Mg , ^{47}Ti and ^{49}Ti

Keyword abstract: NUCLEAR REACTIONS ^{25}Mg , $^{47, 49}\text{Ti}(n, \gamma)$, E=slow; measured $E\gamma$, $I\gamma$. ^{26}Mg , $^{48, 50}\text{Ti}$ deduced transitions.

Keynumber: 1966WAZY

Reference: Proc.Intern.Conf.Study of Nucl.Struct.With Neutrons, Antwerp, Belgium (1965), M.N.de Mevergnies, P.Van Assche, J.Vervier, Eds., North-Holland Publishing Co., Amsterdam, p.536 (1966); EANDC-50-S, Paper 99 (1966)

Authors: R.Wagner, W.M.Good, D.Paya

Title: s-Wave Neutron Strength Functions of Isotopes in the 3s-Resonance Region $40 < A < 70$

Keyword abstract: NUCLEAR REACTIONS ^{43}Ca , $^{47, 49}\text{Ti}$, ^{53}Cr , ^{57}Fe , $^{61}\text{Ni}(n, \gamma)$, E=2-60 keV; $\sigma(\text{nt})$ (E). ^{44}Ca , $^{48, 50}\text{Ti}$, ^{54}Cr , ^{58}Fe , ^{62}Ni deduced resonances, level spacings, strength functions.