

```

=====VIRGIN
PROGRAM VIRGIN VIRGIN
VERSION 76-1 (NOVEMBER 1976) VIRGIN
VERSION 84-1 (JUNE 1984) *DOUBLE PRECISION ENERGY VIRGIN
VERSION 86-1 (JANUARY 1986) *FORTRAN-77/H VERSION VIRGIN
VERSION 88-1 (JULY 1988) *OPTION...INTERNALLY DEFINE ALL I/O VIRGIN
FILE NAMES (SEE, SUBROUTINE FILEIO VIRGIN
FOR DETAILS). VIRGIN
*IMPROVED BASED ON USER COMMENTS. VIRGIN
VERSION 89-1 (JANUARY 1989) *PSYCHOANALYZED BY PROGRAM FREUD TO VIRGIN
INSURE PROGRAM WILL NOT DO ANYTHING VIRGIN
CRAZY. VIRGIN
*UPDATED TO USE NEW PROGRAM CONVERT VIRGIN
KEYWORDS. VIRGIN
*ADDED LIVERMORE CIVIC COMPILER VIRGIN
CONVENTIONS. VIRGIN
VERSION 92-1 (JANUARY 1992) *COMPLETE RE-WRITE VIRGIN
*OUTPUT IN PLOTTAB FORMAT VIRGIN
*UP TO 2000 THICKNESSES VIRGIN
*INCREASED INCORE PAGE SIZE TO 6000 VIRGIN
CROSS SECTION POINTS VIRGIN
*ADDED PHOTON CALCULATIONS VIRGIN
*ADDED BLACKBODY SPECTRUM VIRGIN
*ADDED MULTIPLE LAYERS VIRGIN
*ADDED SPATIALLY DEPENDENT DENSITY VIRGIN
*ADDED FORTRAN SAVE OPTION VIRGIN
*COMPLETELY CONSISTENT I/O ROUTINES - VIRGIN
TO MINIMIZE COMPUTER DEPENDENCE. VIRGIN
VERSION 92-2 (MAY 1992) *CORRECTED TO HANDLE MULTIGROUP CROSS VIRGIN
SECTIONS AS INPUT IN ENDF/B FORMAT. VIRGIN
VERSION 96-1 (JANUARY 1996) *COMPLETE RE-WRITE VIRGIN
*IMPROVED COMPUTER INDEPENDENCE VIRGIN
*ALL DOUBLE PRECISION VIRGIN
*ON SCREEN OUTPUT VIRGIN
*UNIFORM TREATMENT OF ENDF/B I/O VIRGIN
*IMPROVED OUTPUT PRECISION VIRGIN
*DEFINED SCRATCH FILE NAMES VIRGIN
VERSION 99-1 (MARCH 1999) *CORRECTED CHARACTER TO FLOATING VIRGIN
POINT READ FOR MORE DIGITS VIRGIN
*UPDATED TEST FOR ENDF/B FORMAT VIRGIN
VERSION BASED ON RECENT FORMAT CHANGEVIRGIN
*GENERAL IMPROVEMENTS BASED ON VIRGIN
USER FEEDBACK VIRGIN
VERS. 2000-1 (FEBRUARY 2000) *GENERAL IMPROVEMENTS BASED ON VIRGIN
USER FEEDBACK VIRGIN
VERS. 2002-1 (MAY 2002) *OPTIONAL INPUT PARAMETERS VIRGIN
VERS. 2004-1 (MARCH 2004) *ADDED INCLUDE FOR COMMON VIRGIN
*UP TO 2000 THICKNESSES VIRGIN
*INCREASED INCORE PAGE SIZE TO 60,000 VIRGIN
VERS. 2007-1 (JAN. 2007) *CHECKED AGAINST ALL ENDF/B-VII. VIRGIN
*INCREASED INCORE PAGE SIZE TO VIRGIN
240,000 FROM 60,000. VIRGIN
VERS. 2007-2 (DEC. 2007) *72 CHARACTER FILE NAME. VIRGIN
VERS. 2010-1 (Apr. 2010) *General update based on user feedbackVIRGIN
*INCREASED INCORE PAGE SIZE TO VIRGIN
600,000 FROM 240,000. VIRGIN
VERS. 2012-1 (Aug. 2012) *Added CODENAME VIRGIN
*32 and 64 bit Compatible VIRGIN
*Added ERROR stop VIRGIN
VERS. 2015-1 (Jan. 2015) *Extended OUT9. VIRGIN
*Replaced ALL 3 way IF Statements. VIRGIN
*Generalized TART Group Structures. VIRGIN
*Generalized SAND-II Group Structures.VIRGIN
*Extended SAND-II to 60, 150, 200 MeV.VIRGIN
VERS. 2015-2 (Apr. 2015) *Changed ALL data to "D" instead of VIRGIN
"E" to insure it is REAL*8 and avoid VIRGIN
Truncation ERRORS. VIRGIN
VERS. 2017-1 (May 2017) *Added UKAEA 1102 Group Structure. VIRGIN
*Increased points to 3,000,000 VIRGIN
*Increased groupd to 30,000 VIRGIN

```

	*Updated based on user feedback	VIRGIN
	*Defintion of built-in group structure using SUBROUTINE GROPE is identical for GROUPIE and VIRGIN.	VIRGIN
	*All floating point parameters changed to character inout + IN9 conversion.	VIRGIN
VERS. 2018-1 (Jan. 2018)	*Decreased PAGE size from 3,000,000 to 1,500,000	VIRGIN
	*On-line output for ALL ENDERROR	VIRGIN
VERS. 2019-1 (June 2019)	*Additional Interpolation Law Tests	VIRGIN
	*Checked Maximum Tabulated Energy to insure it is the same for all MTs - if not, print WARNING messages.	VIRGIN
VERS. 2020-1 (Feb. 2020)	*Identical to 2019-1.	VIRGIN
VERS. 2023-1 (Feb. 2023)	*Decrease page size from 1,500,000 TO 120,000	VIRGIN

2015-2 Acknowledgment

=====

I thank Andrej Trkov (NDS, IAEA) for finding the problem with the "E" formatted DATA (this effected both VIRGIN and GROUPIE). I also thank Andrej for overseeing the entire PREPRO project at IAEA, Vienna; he is part of a truly International team who worked together to produce PREPRO-2015-2, and to make it available Internationally on-line for FREE to ALL users.

OWNED, MAINTAINED AND DISTRIBUTED BY

THE NUCLEAR DATA SECTION
INTERNATIONAL ATOMIC ENERGY AGENCY
P.O. BOX 100
A-1400, VIENNA, AUSTRIA
EUROPE

ORIGINALLY WRITTEN BY

Dermott E. Cullen

PRESENT CONTACT INFORMATION

Dermott E. Cullen
1466 Hudson Way
Livermore, CA 94550
U.S.A.
Telephone 925-443-1911
E. Mail RedCullen1@Comcast.net
Website RedCullen1.net/HOMEPAGE.NEW

PURPOSE

THIS PROGRAM IS DESIGNED TO CALCULATE UNCOLLIDED (I.E. VIRGIN) FLUX AND REACTIONS DUE TO TRANSMISSION OF A MONODIRECTIONAL BEAM OF NEUTRONS THROUGH ANY THICKNESS OF MATERIAL. IN ORDER TO SIMULATE AN EXPERIMENTAL MEASUREMENT THE RESULTS ARE GIVEN AS INTEGRALS OVER ENERGY TALLY GROUPS (AS OPPOSED TO POINTWISE IN ENERGY). BY TAKING THE RATIO OF REACTIONS TO FLUX IN EACH GROUP AN EQUIVALENT SPATIALLY DEPENDENT GROUP AVERAGED CROSS SECTION IS CALCULATED BY THE PROGRAM.

EVALUATED DATA

THE EVALUATED DATA MUST BE IN THE ENDF/B FORMAT. HOWEVER IT MUST BE LINEAR-LINEAR INTERPOLABLE IN ENERGY-CROSS SECTION BETWEEN TABULATED POINTS. SINCE ONLY CROSS SECTIONS (FILE 3 OR 23) ARE USED, THIS PROGRAM WILL WORK ON ANY VERSION OF ENDF/B (I.E. ENDF/B-I, II, III, IV, V OR VI).

RELATED COMPUTER CODES

IN ORDER TO CONVERT ENDF/B DATA TO THE FORM REQUIRED BY THIS CODE THE FOLLOWING COMPUTER CODES MAY BE USED,

LINEAR - CONVERT FROM GENERAL ENDF/B INTERPOLATION TO LINEAR-LINEAR INTERPOLATION. VIRGIN
 RECENT - ADD THE RESONANCE CONTRIBUTION TO TABULATED BACKGROUND CROSS SECTIONS TO OBTAIN LINEAR-LINEAR INTERPOLABLE RESULTS. VIRGIN
 SIGMA1 - DOPPLER BROADEN CROSS SECTION TO OBTAIN LINEAR-LINEAR INTERPOLABLE RESULTS. VIRGIN
 MIXER - MIX INDIVIDUAL MATERIALS TOGETHER TO DEFINE COMPOSITE MIXTURES, E.G., COMBINE MATERIALS TO DEFINE STAINLESS STEEL. VIRGIN

IN ORDER TO PLOT THE OUTPUT RESULTS OF THIS CODE USE PROGRAM PLOTTAB. VIRGIN

COPIES OF ANY OR ALL OF THESE CODES MAY BE OBTAINED FROM D.E. CULLEN AT THE ABOVE ADDRESS. VIRGIN

OUTPUT FORMAT

 FOR ALL VERSIONS OF THIS PROGRAM PRIOR TO VERSION 92-1 OUTPUT WAS IN TABULAR FORM. VIRGIN
 FOR VERSION 92-1 AND LATER VERSIONS OF THIS CODE ALL OUTPUT IS IN THE PROGRAM PLOTTAB FORMAT TO ALLOW RESULTS TO BE EASILY PLOTTED. FOR A COPY OF PROGRAM PLOTTAB CONTACT D.E. CULLEN AT THE ABOVE ADDRESS. VIRGIN

TALLY GROUPS

 THE TALLY GROUP STRUCTURE MAY BE ANY SET OF MONOTONICALLY INCREASING ENERGY BOUNDARIES. THERE MAY BE UP TO 2000 TALLY GROUPS. BY USING THE INPUT PARAMETERS THE USER MAY SPECIFY ANY ARBITRARY TALLY GROUP STRUCTURE OR SELECT ONE OF THE FOLLOWING BUILT-IN GROUP STRUCTURES. VIRGIN

- (0) TART 175 GROUPS
- (1) ORNL 50 GROUPS
- (2) ORNL 126 GROUPS
- (3) ORNL 171 GROUPS
- (4) SAND-II 620 GROUPS - 1.0D-4 eV UP TO 18 MEV
- (5) SAND-II 640 GROUPS - 1.0D-4 eV UP TO 20 MEV
- (6) WIMS 69 GROUPS
- (7) GAM-I 68 GROUPS
- (8) GAM-II 99 GROUPS
- (9) MUFT 54 GROUPS
- (10) ABBN 28 GROUPS
- (11) TART 616 GROUPS TO 20 MeV
- (12) TART 700 GROUPS TO 1 GeV
- (13) SAND-II 665 GROUPS - 1.0D-5 eV UP TO 18 MEV
- (14) SAND-II 685 GROUPS - 1.0D-5 eV UP TO 20 MEV
- (15) TART 666 GROUPS TO 200 MeV
- (16) SAND-II 725 GROUPS - 1.0D-5 eV UP TO 60 MEV
- (17) SAND-II 755 GROUPS - 1.0D-5 eV UP TO 150 MEV
- (18) SAND-II 765 GROUPS - 1.0D-5 eV UP TO 200 MEV
- (19) UKAEA 1102 GROUPS - 1.0D-5 eV UP TO 1 GeV

INCIDENT SPECTRUM

 THE INCIDENT SPECTRUM MAY BE ANY TABULATED FUNCTION THAT IS GIVEN BY A SET OF POINTS THAT IS MONOTONICALLY INCREASING IN ENERGY AND LINEAR-LINEAR INTERPOLABLE IN ENERGY-SPECTRUM BETWEEN TABULATED POINTS. THERE IS NO LIMIT TO THE NUMBER OF POINTS USED TO DESCRIBE THE SPECTRUM. THERE ARE FIVE BUILT-IN OPTIONS FOR THE SPECTRUM. VIRGIN

- (1) CONSTANT...ENERGY INDEPENDENT (INPUT 0)
- (2) 1/E (INPUT 1)
- (3) BLACKBODY - PHOTON SPECTRUM
- (4) BLACKBODY - ENERGY SPECTRUM (E TIMES THE PHOTON SPECTRUM)
- (5) TRANSMITTED SPECTRUM FROM PREVIOUS CASE

AND AS SUCH THE CYCLE OF TRANSMISSION THROUGH EACH LAYER AND
USING THE TRANSMITTED SPECTRUM AS THE INCIDENT SPECTRUM FOR THE
NEXT LAYER MAY BE REPEATED ANY NUMBER OF TIMES.

REMEMBER - THE INCIDENT SPECTRUM IS ASSUMED TO BE LINEARLY
INTERPOLABLE IN ENERGY AND SPECTRUM BETWEEN THE ENERGIES AT
WHICH IT IS TABULATED. THE TRANSMITTED SPECTRUM WILL BE TABULATED
AT THE UNION OF ALL ENERGIES OF THE INCIDENT SPECTRUM AND CROSS
SECTIONS (TOTAL AND REACTION). IN ORDER TO INSURE THE ACCURACY
OF THE RESULT WHEN PERFORMING MULTIPLE LAYER CALCULATION BE SURE
TO SPECIFY THE INCIDENT SPECTRUM ON THE FIRST LAYER TO SUFFICIENT
DETAIL (ENOUGH ENERGY POINTS CLOSELY SPACED TOGETHER) IN ORDER TO
ALLOW THE TRANSMITTED SPECTRUM TO BE ACCURATELY REPRESENTED BY
LINEAR INTERPOLATION BETWEEN SUCCESSIVE ENERGY POINTS - THERE IS
NO LIMIT TO THE NUMBER OF POINTS ALLOWED IN THE INCIDENT SPECTRUM,
SO IF YOU ARE IN DOUBT, SIMPLY USE MORE ENERGY POINTS TO SPECIFY
THE INCIDENT SPECTRUM.

RESULT OUTPUT UNITS

FLUX = EXACTLY AS CALCULATED
REACTIONS = 1/CM OR 1/GRAM
AVERAGE = 1/CM - MACROSCOPIC UNITS
CROSS SECTION

THICKNESS AND DENSITY

THE UNCOLLIDED CALCULATION ONLY DEPENDS ON THE PRODUCT OF
THICKNESS AND DENSITY (I.E. GRAMS PER CM SQUARED). THIS FACT
MAY BE USED TO SIMPLIFY INPUT BY ALLOWING THE THICKNESS AND
DENSITY TO BE GIVEN EITHER AS CM AND GRAMS/CC RESPECTIVELY
OR ELSE TO GIVE THICKNESS IN GRAMS/(CM*CM) AND INPUT A
DENSITY OF 1.0 - OR IN ANY OTHER CONVENIENT UNITS AS LONG AS
THE PRODUCT OF THICKNESS AND DENSITY IS IN THE CORRECT GRAMS
PER CENTIMETER SQUARED.

GRAMS/(CM*CM) ARE RELATED TO ATOMS/BARN THROUGH THE RELATIONSHIP

$$\text{GRAMS}/(\text{CM}^2) = (\text{ATOMS}/\text{BARN}) * (\text{GRAMS}/\text{MOLE}) * (\text{MOLE}/\text{ATOM})$$

OR...

$$\text{GRAMS}/(\text{CM}^2) = (\text{ATOMS}/\text{BARN}) * (\text{ATOMIC WEIGHT}) / 0.602$$

CROSS SECTIONS AT A SPACE POINT AND OPTICAL THICKNESS

THIS PROGRAM ALLOWS LAYERS OF EITHER UNIFORM DENSITY OR
CONTINUOUSLY VARYING DENSITY. THE DENSITY CAN BE ONE OF THE
FOLLOWING FORMS,

- | | | |
|-----------------------|--------------------------------|--------|
| 1) C | = UNIFORM DENSITY | VIRGIN |
| 2) C*2*(X/T) | = LINEAR VARIATION FROM 0 TO C | VIRGIN |
| 3) C*(2-2*(X/T)) | = LINEAR VARIATION FROM C TO 0 | VIRGIN |
| 4) C*3*(X/T)**2 | = SQUARE VARIATION FROM 0 TO C | VIRGIN |
| 5) C*(3-3*(X/T)**2)/2 | = SQUARE VARIATION FROM C TO 0 | VIRGIN |
| 6) C*4*(X/T)**3 | = CUBIC VARIATION FROM 0 TO C | VIRGIN |
| 7) C*(4-4*(X/T)**3)/3 | = CUBIC VARIATION FROM C TO 0 | VIRGIN |

IN ORDER TO CALCULATE REACTIONS AT A POINT THE MICROSCOPIC
REACTION CROSS SECTION NEED MERELY BE SCALED BY THESE DENSITIES.

IN ORDER TO CALCULATE TRANSMISSION WE MUST DEFINE THE OPTICAL
PATH LENGTH WHICH MAY BE DEFINED BY INTEGRATING EACH OF THE
ABOVE DENSITY FORMS TO FIND,

- | | |
|-----------------------|--------|
| 1) C*X | VIRGIN |
| 2) C*X*(X/T) | VIRGIN |
| 3) C*X*(2-(X/T)) | VIRGIN |
| 4) C*X*(X/T)**2 | VIRGIN |
| 5) C*X*(3-(X/T)**2)/2 | VIRGIN |
| 6) C*X*(X/T)**3 | VIRGIN |
| 7) C*X*(4-(X/T)**3)/3 | VIRGIN |


```

F(X)=0.5*(FK*(1 - X) + FK+1*(1 + X))
      =0.5*(FK+1 + FK) + 0.5*(FK+1 - FK)*X
VIRGIN
VIRGIN
VIRGIN
BY DEFINING THE AVERAGE VALUE AND 1/2 THE CHANGE ACROSS THE
INTERVAL.
VIRGIN
VIRGIN
VIRGIN
VIRGIN
VIRGIN
AVF=0.5*(FK+1 + FK)
DF= 0.5*(FK+1 - FK)
DE= 0.5*(EK+1 - EK)
VIRGIN
VIRGIN
VIRGIN
EACH OF THE THREE FUNCTIONS REDUCES TO THE SIMPLE FORM...
VIRGIN
VIRGIN
VIRGIN
F(X)=AVF+DF*X
VIRGIN
VIRGIN
AND THE TWO REQUIRED INTEGRALS REDUCE TO...
VIRGIN
VIRGIN
VIRGIN
FLUX
----
DE*EXP(-AVXCT*Z) * (INTEGRAL -1 TO +1)
((AVS+DS*X)*EXP(-DXCT*Z*X)*DX)
VIRGIN
VIRGIN
VIRGIN
REACTION
-----
DE*EXP(-AVXCT*Z) * (INTEGRAL -1 TO +1)
((AVS*AVXCR+(AVS*DXCR+AVXCR*DS)*X+DS*DXCR*X*X)*EXP(-DXCT*Z*X)*DX)
VIRGIN
VIRGIN
VIRGIN
WHERE
VIRGIN
VIRGIN
VIRGIN
AVXCT = AVERAGE VALUE OF THE TOTAL CROSS SECTION
AVXCR = AVERAGE VALUE OF THE REACTION CROSS SECTION
AVS   = AVERAGE VALUE OF THE SOURCE
DXCT  = 1/2 THE CHANGE IN THE TOTAL CROSS SECTION
DXCR  = 1/2 THE CHANGE IN THE REACTION CROSS SECTION
DS    = 1/2 THE CHANGE IN THE SOURCE
DE    = 1/2 THE CHANGE IN THE ENERGY
VIRGIN
VIRGIN
VIRGIN
NOTE THAT IN THIS FORM THE ENERGY ONLY APPEARS IN FRONT OF THE
INTEGRALS AND THE INTEGRALS ARE EXPRESSED ONLY IN TERMS OF THE
TABULATED VALUES OF S(E), XCT(E) AND XCR(E). IN PARTICULAR NO
DERIVATIVES ARE USED, SO THAT THERE ARE NO NUMERICAL INSTABILITY
PROBLEMS IN THE VICINITY OF DISCONTINUITIES IN S(E), XCT(E) OR
XCR(E). INDEED, SINCE (EK+1 - EK) APPEARS IN FRONT OF THE INTEGRAL
POINTS OF DISCONTINUITY AUTOMATICALLY MAKE ZERO CONTRIBUTION TO
THE INTEGRALS.
VIRGIN
VIRGIN
VIRGIN
THE REQUIRED INTEGRALS CAN BE EXPRESSED IN TERMS OF THE THREE
INTEGRALS IN NORMAL FORM...
VIRGIN
VIRGIN
VIRGIN
F(A,N) = (INTEGRAL -1 TO 1) (X**N*EXP(-A*X)*DX), N=0,1 AND 2.
VIRGIN
VIRGIN
VIRGIN
THESE THREE INTEGRALS CAN BE EVALUATED TO FIND...
VIRGIN
VIRGIN
VIRGIN
N=0
---
F(A,0) = (EXP(A)-EXP(-A))/A
VIRGIN
VIRGIN
VIRGIN
N=1
---
F(A,1) = ((1-A)*EXP(A)-(1+A)*EXP(-A))/(A*A)
VIRGIN
VIRGIN
VIRGIN
N=2
---
F(A,2) = ((2-2*A+A*A)*EXP(A)-(2+2*A+A*A)*EXP(-A))/(A*A*A)
VIRGIN
VIRGIN
VIRGIN
HOWEVER THESE EXPRESSIONS ARE NUMERICALLY UNSTABLE FOR SMALL
VALUES OF A. THEREFORE FOR SMALL A THE EXPONENTIAL IN THE
INTEGRALS ARE EXPANDED IN A POWER SERIES...
VIRGIN
VIRGIN
VIRGIN
EXP(-AX)=1.0-(AX)+(AX)**2/2-(AX)**3/6+(AX)**4/24-.....
=(SUM K=0 TO INFINITY) (-AX)**K/(K FACTORIAL)
VIRGIN
VIRGIN

```

AND THE INTEGRAL REDUCES TO THE FORM...

$$(SUM K=0 TO INFINITY) ((-A)**K/(K FACTORIAL)) * (INTEGRAL -1 TO 1) (X**(N+K))*DX$$

WHICH CAN BE ANALYTICALLY EVALUATED TO FIND...

(K(N) = K FACTORIAL)

N=0

F(A,0) = 2*(1+(A**2)/K(3)+(A**4)/K(5)+(A**6)/K(7)+....

N=1

F(A,1) = -2*A*(2/K(3)+4*(A**2)/K(5)+6*(A**4)/K(7)+8*(A**6)/K(9)+..

N=2

F(A,2) = 2*(2/K(3)+3*4*(A**2)/K(5)+5*6*(A**4)/K(7)+7*8*(A**6)/K(9)+....

THESE EXPANSIONS ARE USED WHEN THE ABSOLUTE VALUE OF A IS LESS THAN 0.1. BY TRUNCATING THE ABOVE SERIES BEFORE A**8 THE ERROR RELATIVE TO THE LEADING TERM OF THE SERIES WILL BE 10**(-10), YIELDING 10 DIGIT ACCURACY.

AFTER EVALUATING THE ABOVE FUNCTIONS, EITHER DIRECTLY OR BY USING THE EXPANSION THE TWO REQUIRED INTEGRALS CAN BE WRITTEN AS...

FLUX

DE*EXP(-AVXCT*Z)*(AVS*F(A,0) + DS*F(A,1))

REACTIONS

DE*EXP(-AVXCT*Z)*
(AVS*AVXCR*F(A,0) + (AVS*DXCR+AVXCR*DS)*F(A,1) + DS*DXCR*F(A,2))

INPUT FILES

FILENAME UNIT DESCRIPTION

INPUT 2 INPUT LINES
ENDFIN 10 EVALUATED DATA IN ENDF/B FORMAT

OUTPUT FILES

FILENAME UNIT DESCRIPTION

OUTPUT 3 OUTPUT LISTING

SCRATCH FILES

FILENAME UNIT DESCRIPTION

SCR1 12 REACTION, FLUX AND CROSS SECTION RESULTS (BCD)
(SORTED AT END OF RUN AND OUTPUT SEPARATELY)
SCR2 13 TALLY GROUP ENERGY BOUNDARIES (BINARY)
SCR3 14 SOURCE SPECTRUM (BINARY)
SCR4 15 TOTAL CROSS SECTION (BINARY)
SCR5 16 REACTION CROSS SECTION (BINARY)

OPTIONAL STANDARD FILE NAMES (SEE SUBROUTINE FILIO1 AND FILEIO2)

UNIT FILE NAME FORMAT

2 VIRGIN.INP BCD
3 VIRGIN.LST BCD
10 ENDFB.IN BCD
11-15 (SCRATCH) BINARY
16 PLOTTAB.CUR PLOTTAB OUTPUT FORMAT DATA

LINE	COLS.	FORMAT	DESCRIPTION	
INPUT LINES				VIRGIN
-----				VIRGIN
ANY NUMBER OF CASES MAY BE RUN ONE AFTER THE OTHER. AFTER THE				VIRGIN
FIRST CASE HAS BEEN RUN THE FOLLOWING CASES MAY USE THE SAME				VIRGIN
THICKNESSES, GROUP STRUCTURE AND SPECTRUM AS THE PRECEDING CASE.				VIRGIN
IN ADDITION THE TRANSMITTED SPECTRUM FROM ONE CASE MAY BE USED				VIRGIN
AS THE INCIDENT SPECTRUM IN THE NEXT CASE, TO ALLOW MULTIPLE				VIRGIN
LAYERS OF DIFFERENT MATERIALS.				VIRGIN
				VIRGIN
1	1-60	ENDF/B	INPUT DATA FILENAME (STANDARD OPTION = ENDFB.IN)	VIRGIN
				VIRGIN
LEAVE THE DEFINITION OF THE FILENAMES BLANK - THE PROGRAM WILL				VIRGIN
THEN USE STANDARD FILENAMES.				VIRGIN
				VIRGIN
2-3	1-72	18A4	TWO LINE TITLE DESCRIBING PROBLEM	VIRGIN
4	1- 6	I6	ZA (1000*Z+A) OF TARGET FOR TOTAL	VIRGIN
	7-11	I5	MT OF TOTAL	VIRGIN
	12-22	E11.4	DENSITY FOR TOTAL	VIRGIN
	23-28	I6	ZA (1000*Z+A) OF TARGET FOR REACTION	VIRGIN
	29-33	I5	MT OF REACTION	VIRGIN
			= 0 - NO REACTION CALCULATION (ONLY FLUX).	VIRGIN
			= GREATER THAN 0 - CALCULATE REACTIONS.	VIRGIN
34-44	E11.4		DENSITY FOR REACTION	VIRGIN
45-50	I6		NUMBER OF TARGET THICKNESSES	VIRGIN
			= GREATER THAN 0 = READ FROM INPUT	VIRGIN
			(1 TO 2000 ALLOWED)	VIRGIN
			= 0 = SAME AS LAST CASE	VIRGIN
51-55	I5		NUMBER OF TALLY GROUPS	VIRGIN
			(REMEMBER NUMBER OF GROUP BOUNDARIES	VIRGIN
			IS ONE MORE THAN THE NUMBER OF GROUPS)	VIRGIN
			UP TO 2000 GROUPS ARE ALLOWED	VIRGIN
			BUILT-IN GROUP STRUCTURES.	VIRGIN
			= GREATER THAN 0 = READ FROM INPUT	VIRGIN
			= 0 TART 175 GROUPS	VIRGIN
			= -1 ORNL 50 GROUPS	VIRGIN
			= -2 ORNL 126 GROUPS	VIRGIN
			= -3 ORNL 171 GROUPS	VIRGIN
			= -4 SAND-II 620 GROUPS..1.0D-4 eV TO 18 MEV	VIRGIN
			= -5 SAND-II 640 GROUPS..1.0D-4 eV TO 20 MEV	VIRGIN
			= -6 WIMS 69 GROUPS	VIRGIN
			= -7 GAM-I 68 GROUPS	VIRGIN
			= -8 GAM-II 99 GROUPS	VIRGIN
			= -9 MUFT 54 GROUPS	VIRGIN
			=-10 ABBN 28 GROUPS	VIRGIN
			=-11 TART 616 GROUPS TO 20 MeV	VIRGIN
			=-12 TART 700 GROUPS TO 1 GeV	VIRGIN
			=-13 SAND-II 665 GROUPS..1.0D-5 eV TO 18 MEV	VIRGIN
			=-14 SAND-II 685 GROUPS..1.0D-5 eV TO 20 MEV	VIRGIN
			=-15 TART 666 GROUPS TO 200 MeV	VIRGIN
			=-16 SAND-II 725 GROUPS..1.0D-5 eV TO 60 MEV	VIRGIN
			=-17 SAND-II 755 GROUPS..1.0D-5 eV TO 150 MEV	VIRGIN
			=-18 SAND-II 765 GROUPS..1.0D-5 eV TO 200 MEV	VIRGIN
			=-19 UKAEA 1102 GROUPS..1.0D-5 eV to 1 GeV	VIRGIN
56-60	I5		NUMBER OF POINTS IN SOURCE SPECTRUM	VIRGIN
			(MUST BE AT LEAST TWO POINTS)	VIRGIN
			= GREATER THAN 1 = READ FROM INPUT	VIRGIN
			= 0 = SAME AS LAST CASE	VIRGIN
			= -1 = CONSTANT (ENERGY INDEPENDENT)	VIRGIN
			= -2 = 1/E	VIRGIN
			= -3 = BLACKBODY - PHOTON SPECTRUM	VIRGIN
			= -4 = BLACKBODY - ENERGY SPECTRUM	VIRGIN
			= -5 = TRANSMITTED SPECTRUM FROM LAST CASE	VIRGIN
			NOTE, ALL SPECTRA, EXCEPT THE TRANSMITTED	VIRGIN
			SPECTRUM FROM THE LAST CASE, WILL BE	VIRGIN
			NORMALIZED SUCH THAT ITS INTEGRAL OVER	VIRGIN
			ENERGY WILL BE UNITY.	VIRGIN
61-64	1X,3I1		SPATIALLY DEPENDENT OUTOUT	VIRGIN

			= 0 = NO	VIRGIN
			= 1 = YES	VIRGIN
			FOR THE 3 QUANTITIES	VIRGIN
			COLUMN 67 FLUX	VIRGIN
			68 REACTIONS	VIRGIN
			69 AVERAGE CROSS SECTION	VIRGIN
65-65	I1		ENERGY DEPENDENT OUTOUT	VIRGIN
			= 0 = NONE	VIRGIN
			= 1 = INCIDENT SPECTRUM	VIRGIN
			= 2 = TRANSMITTED SPECTRUM	VIRGIN
			= 3 = INCIDENT REACTIONS	VIRGIN
			= 4 = TRANSMITTED REACTIONS	VIRGIN
			= 5 = TOTAL CROSS SECTION	VIRGIN
			= 6 = REACTION CROSS SECTION	VIRGIN
5	1-11	E11.4	BLACKBODY TEMPERATURE IN eV	VIRGIN
	12-22	E11.4	FLUX NORMALIZATION	VIRGIN
	23-33	E11.4	REACTION NORMALIZATION	VIRGIN
			CALCULATIONS WILL BE BASED ON THE SPECTRUM	VIRGIN
			AND CROSS SECTIONS AS READ. AT OUTPUT THE	VIRGIN
			RESULTS WILL BE MULTIPLIED BY THESE	VIRGIN
			NORMALIZATION FACTORS.	VIRGIN
	34-44	I11	DENSITY PROFILE	VIRGIN
			= 0 - UNIFORM - BASED ON TOTAL DENSITY	VIRGIN
			= 1 - UNIFORM - TOTAL + REACTION DENSITY	VIRGIN
			= 2 - TOTAL + LINEAR REACTION	VIRGIN
			= 3 - LINEAR (TOTAL + REACTION)	VIRGIN
			= 4 - TOTAL + SQUARE REACTION	VIRGIN
			= 5 - SQUARE (TOTAL + REACTION)	VIRGIN
			= 6 - TOTAL + CUBIC REACTION	VIRGIN
			= 7 - CUBIC (TOTAL + REACTION)	VIRGIN
6-N	1-66	6E11.4	TARGET THICKNESSES IN CM	VIRGIN
			IF SAME AS LAST CASE THIS SECTION IS NOT	VIRGIN
			INCLUDED IN THE INPUT.	VIRGIN
VARY	1-66	6E11.4	TALLY GROUP ENERGY BOUNDARIES	VIRGIN
			(NUMBER OF BOUNDARIES IS ONE MORE THAN	VIRGIN
			THE NUMBER OF TALLY GROUPS)	VIRGIN
			IF THE STANDARD OPTION (-14 TO 0) IS	VIRGIN
			SELECTED THIS SECTION IS NOT INCLUDED	VIRGIN
			IN THE INPUT	VIRGIN
VARY	1-66	6E11.4	SOURCE SPECTRUM IN ENERGY (eV)-SOURCE PAIRS	VIRGIN
			(MUST BE AT LEAST TWO POINTS)	VIRGIN
			IF STANDARD OPTION (-5 TO 0) IS SELECTED THIS	VIRGIN
			SECTION IS NOT INCLUDED IN THE INPUT	VIRGIN
				VIRGIN
			ANY NUMBER OF CASES MAY BE RUN ONE AFTER ANOTHER.	VIRGIN
				VIRGIN
			EXAMPLE INPUT NO. 1	VIRGIN
			-----	VIRGIN
			CALCULATE THE UNCOLLIDED FLUX AND CAPTURE (MT=102) THROUGH	VIRGIN
			30 CM OF IRON (DENSITY 7.87 G/CC). TALLY THE RESULTS USING	VIRGIN
			THE TART 175 GROUP STRUCTURE. THE SOURCE WILL BE CONSTANT	VIRGIN
			FROM 1 KEV TO 20 MEV. USE THE STANDARD ENDF/B INPUT DATA	VIRGIN
			FILENAME.	VIRGIN
				VIRGIN
			ENDFB.IN	VIRGIN
			IRON 0 TO 30 CM THICK.	VIRGIN
			CONSTANT SOURCE FROM 1 KEV TO 20 MEV.	VIRGIN
			26000 1 7.8700D+ 0 26000 102 7.8700D+ 0 2 0 2 1100	VIRGIN
			0.0000D+ 0 1.0000D+ 0 1.0000D+ 0 0 0.0000D+00	VIRGIN
			0.0000D+00 3.0000D+01	VIRGIN
			1.0000D+03 1.0000D+00 2.0000D+07 1.0000D+00	VIRGIN
				VIRGIN
			EXAMPLE INPUT NO. 2	VIRGIN
			-----	VIRGIN
			CALCULATE THE UNCOLLIDED PHOTON FLUX THROUGH A MIXTURE OF SILICON	VIRGIN
			AND IRON FOR 100 MEV PHOTONS INCIDENT. THE TRANSMISSION WILL BE	VIRGIN
			CALCULATED FOR 21 THICKNESSES VARYING BETWEEN 0 AND 1 CM. THERE	VIRGIN
			WILL BE ONLY 1 TALLY GROUP SPANNING A VERY NARROW ENERGY RANGE	VIRGIN
			NEAR 100 MEV, AND THE SOURCE SPECTRUM WILL BE CONSTANT OVER THE	VIRGIN
			SAME ENERGY RANGE. USE THE STANDARD ENDF/B INPUT DATA FILENAME	VIRGIN
			BY LEAVING THE FIRST INPUT LINE BLANK.	VIRGIN

```
(THIS IS A BLANK LINE TO USE THE STANDARD INPUT FILENAME) VIRGIN
100 MEV PHOTONS VIRGIN
SILICON + 5 % IRON VIRGIN
14000 521 2.30000+ 0 26000 521 1.15000- 1 21 1 2 1000 VIRGIN
0.00000+ 0 1.00000+ 0 1.00000+ 0 1 0.00000+00 VIRGIN
0.00000+00 5.00000-01 1.00000+00 1.50000+00 2.00000+00 2.50000+00VIRGIN
3.00000+00 3.50000+00 4.00000+00 4.50000+00 5.00000+00 5.50000+00VIRGIN
6.00000+00 6.50000+00 7.00000+00 7.50000+00 8.00000+00 8.50000+00VIRGIN
9.00000+00 9.50000+00 1.00000+01 VIRGIN
9.99000+ 7 1.00100+ 8 VIRGIN
9.99000+ 7 1.00000+ 4 1.00100+ 8 1.00000+ 4 VIRGIN
=====VIRGIN
```