




XA9745762



International Atomic Energy Agency

INDC(HUN)-032

Distr. L

IN DC

INTERNATIONAL NUCLEAR DATA COMMITTEE

**INVESTIGATIONS ON (n,p) CROSS SECTIONS
IN THE 14 MeV REGION**

R. Dóczki¹, V. Semkova², A.D. Majdeddin³, Cs.M. Buczkó¹ and J. Csikai¹

¹Institute of Experimental Physics, Kossuth University
H-4001 Debrecen, Pf. 105, Hungary

²Institute of Nuclear Physics and Nuclear Energy, Bulgarian Academy of Sciences
Tzarigradsko chausse 72, 1784 Sofia, Bulgaria

³Faculty of Nuclear and Electronic Engineering, Al-Fateh University
P.O. Box 13292, Tripoli, Libya

October 1997

IAEA NUCLEAR DATA SECTION, WAGRAMERSTRASSE 5, A-1400 VIENNA

29 - 02 -

Reproduced by the IAEA in Austria
October 1997

INDC(HUN)-032

Distr. L

**INVESTIGATIONS ON (n,p) CROSS SECTIONS
IN THE 14 MeV REGION**

R. Dóczi¹, V. Semkova², A.D. Majdeddin³, Cs.M. Buczkó¹ and J. Csikai¹

**¹Institute of Experimental Physics, Kossuth University
H-4001 Debrecen, Pf. 105, Hungary**

**²Institute of Nuclear Physics and Nuclear Energy, Bulgarian Academy of Sciences
Tzarigradsko chausse 72, 1784 Sofia, Bulgaria**

**³Faculty of Nuclear and Electronic Engineering, Al-Fateh University
P.O. Box 13292, Tripoli, Libya**

October 1997

INVESTIGATIONS ON (n,p) CROSS SECTIONS IN THE 14 MeV REGION

R. Dóczl¹, V. Semkova², A. D. Majdeddin³, Cs. M. Buczkó¹ and J. Csikai¹*

¹ Institute of Experimental Physics, Kossuth University
H-4001 Debrecen, Pf. 105., Hungary

²Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences
Tzarigradsko chausse 72, 1784 Sofia, Bulgaria

³ Faculty of Nuclear and Electronic Engineering
Al-Fateh University P. O. Box 13292, Tripoli, Libya

Cross sections have been measured, deduced, and adopted for 208 (n,p) reactions at (14.7 ± 0.2) MeV incident neutron energy. Systematics based on asymmetry parameter dependence and isotopic effects have been confirmed. Recommended formulae were fitted to the same data base to be able to test their applicability. Total (n,p) cross sections are given for 54 elements and compared with the results of direct methods. Some $\sigma_{n,p}$ data both for long-lived target and residual radionuclides were estimated.

1. Introduction

Recently, a number of hydrogen production cross section have been measured and calculated for elements and isotopes at around 14 MeV neutron energy [1-14], mainly for the radiation damage assessment of fusion related materials, to complete the data base of neutron activation cross sections for environmental protection, and to estimate the potential radiation hazards connected with the nuclear techniques. Cross sections for the generation of long-lived radionuclides, of importance in fusion reactor technology and accelerator based neutron

* Author for correspondence.

sources, have a high priority. Precise data at 14 MeV are needed for validation and testing of different data libraries and for the normalization of excitation functions.

Recently, on the basis of precise cross section measurements a number of investigations [11,27,81,83-87] were devoted to improve the systematics based on the isotopic and asymmetry parameter dependences. The reliable theoretical and empirical formulae are indispensable to estimate cross sections if experimental data are not available.

2. Experimental procedure

The description of the sample preparation, the properties of neutron sources, the irradiation conditions as well as the absolute activity determination were summarized in our recent paper [15] on the investigations of (n,α) activation cross sections around 14 MeV.

3. Results and conclusions

Activation cross sections of (n,p) reactions, measured and deduced in Debrecen, were completed with the recent literature data in order to prepare a recommended library of the consistent data. A systematic study of the cross sections has shown that some data are discrepant due to the different measuring procedures, the adopted reference reactions and decay parameters. The recommended cross sections are given as the average of the most recent data [8-12, 16-80] by omitting those which show unrealistic deviations from the mean. The recommended (n,p) cross sections are summarized in Table 1.

The experimental data measured around 14 MeV can be extrapolated to a given incident neutron energy by using the following expression [27] for the slopes (m_r) of the excitation functions

$$m_r (\%/\text{MeV}) = -19.1 + 137.8S + 1207S^2 , \quad (1)$$

where $S=(N-Z)/A$ is the asymmetry parameter.

Kasugai et al. [27] have given the empirical formula, Eq. 2, to determine the σ_{14} values in a wide mass number region ($A=19-187$)

$$\sigma_{14} (\text{mb}) = 1830(N-Z+1) \exp(-50.7(N-Z+1)/A) . \quad (2)$$

Equations (1) and (2) can be used for the estimation of the cross sections in the 14 MeV region. It was found in Ref. 15, that similarly to the (n,α) reactions, the isotopic dependence of the measured and deduced (n,p) cross sections can be well approximated by the following expression

$$\sigma_{n,p}(A)_{Z=\text{const}} = ae^{-bS-cS^2} , \quad (3)$$

where a , b , and c are fitting parameters. Some examples for the $\sigma_{n,p}(A)$ functions are shown in Fig. 1.

As shown in Fig. 1 the isotopic dependence of $\sigma_{n,p}$ values is not linear in a semi-log plot.

Using Eq. 3 61 unknown $\sigma_{n,p}$ data were deduced for nuclei having unfavourable decay properties. These cross sections are given in Table 2.

Total (n,p) cross sections were determined for 54 elements by averaging the $\sigma_{n,p}$ values over isotopic abundances. These data are summarized in Table 3.

The analyses of the available $\sigma_{n,p-\text{em}}$ values [1-14] and the recommended $\sigma_{n,p}$ data indicate that the (n,np) and (n,d) reactions are significant even for medium and heavy nuclei.

The fit of the empirical expression given by Levkovskii [81,82] to the data summarized in Tables 1 and 2 proved that the gross trend in the (n,p) cross sections is determined by the

asymmetry parameter S. A new formula (Eq. 5), with an additional term in the exponential, gives a substantial improvement in fitting the data [15]

$$\sigma_{n,p} = a(A^{1/3} + 1)^2 e^{-b(S + S^2)} = aB(A)e^{-b(S + S^2)}. \quad (5)$$

The values of a and b parameters are 23.659 ± 1.128 and 23.041 ± 0.507 , respectively.

In order to eliminate the two parameter dependence in Eq. 5, the $\sigma_{n,p}$ values were divided by the absorption term B.

$$R = \sigma_{n,p} / (A^{1/3} + 1)^2 = e^{-bS + cS^2} \quad (6)$$

Fitted R ratios and the values of a, b, and c parameters are given in Fig.2. This figure shows that the deviations from the gross trend are significant for magic Z numbers and deformed nuclei. The fits of the formulae to the data were based on the weighted least-squares method. To judge on the systematics quality the following quantity was determined $F = \sum (|\sigma_{\text{exp}} - \sigma_{\text{calc}}|) / \sigma_{\text{calc}}$. For weighting in F neither σ_{exp} nor $\Delta\sigma$ are recommended.

The quantity F/n , where n is the number of data points available in the intervals covered by the given systematics, can characterize the goodness of the formulae.

As shown in Table 4, the empirical formulae given by Eder et al. [83], Bychkov et al. [84], Forrest [11], Ait-Tahar [85], and Kasugai et al. [27] agree well with the recommended (n,p) cross sections in wide Z and A intervals without splitting the data library. It should be noted, however, that the formulae consisting of different parts for different regions of Z or A give better fits to the data. Histograms in Figs.3a, 3b and 3c show the fit of the different formulae to the (n,p) data given in Tables 1 and 2.

The systematic studies of the $\sigma_{n,p}$ data rendered possible to estimate some (n,p) cross sections both for long-lived target and residual radionuclides. These data are summarized in Table 5.

It should be noted, that the current version of the SINCROS-II system [88] is able to describe the (n,p) cross section curves up to 50 MeV. The SINCROS-II calculations [89-91] agree well with the 14 MeV data.

Acknowledgments

The authors are grateful to Dr. S. Szegedi, Ms. A. Liszka, Mr. D. Bolyán and Mr. J. Szegedi for the technical assistance. Thanks are due to Dr. M. Várnagy for the valuable discussion on the fitting methods.

This work was supported in part by the Hungarian Research Fund (Contract No. T 016713/95), and the International Atomic Energy Agency, Vienna (Contract No. 7687/R3).

References

- [1] R.C.Haight, S.M.Grimes, R.G.Johson and H.H.Barschall, Charged-Particle Emission in Reactions of 15 MeV Neutrons with ^{89}Y , ^{90}Zr and $^{92,94,95,96}\text{Mo}$, Phys. Rev. C23(1981)700-707.
- [2] S.M.Grimes, R.C.Haight, K.R.Alvar, H.H.Barschall and R.R.Borchers, Charged-Particle Emission in Reactions of 15 MeV Neutrons with Isotopes of Chromium, Iron, Nickel and Copper, Phys. Rev.C19(1979)2127-2137.
- [3] R.Fischer, G.Traxler, H.Vonach, and M.Uhl, Investigation of Charged Particles from (n,p) and (n, α) Reactions, Rad. Effects, 96(1986)309-312.
- [4] S.M.Grimes, R.C.Haight and J.D.Anderson, Charged-Particle-Producing Reactions of 15 MeV Neutrons on ^{51}V and ^{93}Nb , Phys.Rev.C17 (1978)508-515.
- [5] S.M.Grimes, R.C. Haight and J.D.Anderson, Measurement of Sub-Coulomb-Barrier Charged Particles Emitted from Aluminium and Titanium Bombarded by 15 MeV Neutrons, Nucl. Sci. Eng.62(1977)187-194.
- [6] L.R.Greenwood, D.G.Doran and H.L.Heinisch, Production of ^{91}Nb , ^{94}Nb and ^{95}Nb from Mo by 14.5-14.8 MeV Neutrons, Phys.Rev.35(1987)76-80.
- [7] C.Tsabarlis, E.Wattecamp and G.Rollin, Double Differential (n,xp) and (n,x α) Cross Section Measurements of ^{27}Al , ^{58}Ni and ^{63}Cu in the Neutron Energy Range from 2.0 to 15.5 MeV, ibid. pp.282-284.

- [8] CINDA-Series: The Index to Literature and Computer Files on Microscopic Neutron Data, IAEA, Vienna, 1990 and 1995.
- [9] V.McLane, C.L.Dunford and P.F.Rose, Neutron Cross Sections, Vol.2, Academic Press, Inc., New York, 1988.
- [10] J.Csikai, Handbook of Fast Neutron Generators, CRC Press.Inc., Boca Raton, Florida, 1987, Vols.I and II.
- [11] R.A.Forrest, Systematics of Neutron-Induced Threshold Reactions with Charged Products at about 14.5 MeV, AERE-R 12419, Harwell Laboratory, (December,1986).
- [12] J.Kopecky, Experimental Data Base Compiled for Renormalizations and Validation of EAF-4.1 Data, EAF-Doc-6,7,8, ECN Petten, (August 1994, September 1994, January1996).
- [13] R. C. Haight, S. M. Grimes, and J. D. Anderson, Hydrogen and Helium Production Cross Sections for 15 MeV Neutrons on Types 316 and 304 Stainless Steel, Nuclear Science and Engineering: 63, 200-204, 1977.
- [14] B. A. Magurno, Dosimetry, Gas Production, and Activation Cross-Section Data, Progress in Nuclear Energy, Vol 13, No 2/3, pp. 259-285, 1984.
- [15] A. D. Majdeddin, V. Semkova, R. Dóczzi, Cs. M. Buczkó and J Csikai, Investigations on (n,α) cross sections in the 14 MeV region, INDC(HUN), IAEA, Vienna (in press).
- [16] V M.Bychkov, V.N.Manokhin, A.B.Pashchenko and V.I.Plyashin, Cross Sections for the (n,p) , (n,α) and $(n,2n)$ Threshold Reactions, INDC(CCP)-146, IAEA, Vienna, 1980.
- [17] A.B.Pashchenko, Reaction Cross-Sections Induced by 14.5 MeV and by Cf-252 and U-235 Fission Spectrum Neutrons, INDC(CCP)-323, IAEA, Vienna, 1991.
- [18] S.M.Qaim, 14 MeV Neutron Activation Cross-Sections, in Handbook of Spectroscopy, Vol.III, CRC Press, Inc., Boca Raton, Florida, 1981, p.141.
- [19] C.Konno, Y.Ikeda, K.Oishi, K.Kawade, H.Yamamoto and H.Maekawa, Activation Cross Section Measurements at Neutron Energy from 13.3 to 14.9 MeV Using the FNS Facility, JAERI 1329, Tokai-mura (October,1993).
- [20] Y.Ikeda, Ch.Konno, K.Oishi, T.Nakamura, H.Miyade, K.Kawade, H.Yamamoto and T.Katoh, Activation Cross Section Measurement for Fusion Reactor Structural

- Materials at Neutron Energy from 13.3 to 15.0 MeV Using FNS Facility, JAERI
1312,Tokai-mura (March,1988).**
- [21] R.C.Ward, I.C.Gomes and D.L.Smith, A Survey of Selected Neutron-Activation Reactions with Short-Lived Products of Importance to Fusion Reactor Technology, INDC(USA)-106, IAEA, Vienna, (1994).
 - [22] T.Katoh, K.Kawade and H.Yamamoto, Measurement of Activation Cross Sections, JAERI-M 89-083, Tokai-mura (1989).
 - [23] N.Yamamuro and S.Iijima, Activation Cross Section Data File(I), JAERI-M 89-129, Tokai-mura(1989).
 - [24] C.B.A.Forty, R.A.Forrest, D.J.Compton and C.Royner, Handbook of Fusion Activation Data, AEA Fus 180(May,1992), AEA Fus 232(May,1993), AEA Technology, Culham Laboratory, Abingdon, UK.
 - [25] N.B.Perez, E.T.Cheng, A.B.Pashchenko and H.K.Vonach, Present Status of Activation Cross Section Data for Production of Long-Lived Radionuclides Probably most Important for Waste Disposal, INDC(NDS)-302, IAEA, Vienna, (1994).
 - [26] A.A.Filatenkov, S.V.Chuvaev, V.N.Absenov and V.A.Jakovlev, Systematic Measurement of Activation Cross Section at Neutron Energies from 13.4-14.9 MeV, INDC(CCP)-402(January,1997).
 - [27] Y.Kasugai, Y.Ikeda, H.Yamamoto and K.Kawade, Systemetics for (n,p) excitation functions in the neutron energy between 13.3 and 15.0 MeV, Ann. Nucl. Energy Vol. 23 No.18, pp1429-1444. (1996).
 - [28] K.Gul, Systematics of (n,p) and (n, α) Cross Sections for 14 MeV Neutrons on Basis of Statistical Model, INDC(PAK)-009 (July,1995), IAEA, Vienna.
 - [29] Nuclear Data For Fusion Reactor Technology, (Ed. S.Cierjacks), NEANDC-307 'U',INDC(GER)-035 (April,1992), KfK, Karlsruhe,5062.
 - [30] Zhao Wenrong, Lu Hanlin, Yu Weixiang and Yuan Xialin, Compilation of Measurements and Evaluations of Nuclear Activation Cross Sections for Nuclear Data Applications, INDC(CPR)-16, (Agust,1989), IAEA, Vienna.
 - [31] H.Liskien, R.Wölfle, R.Widera and S.M.Qaim, Excitation Functions of (n,p) and (n, α) Reactions on Molybdenum Isotopes, Appl. Radiat.Isot. 41(1990)83-90.

- [32] N.I Molla , R.U.Miah, M.Rahman and Aysha Akhter, Excitation Functions of Some (n,p), (n,2n) and (n, α) Reactions on Nickel, Zircunium and Niobium Isotopes in the Energy Range 13.64-14.83 MeV, ibid.pp.355-357.
- [33] K.Kawade, H.Yamamoto, T.Katoh, A.Taniguchi, T.Ikuta, Y.Kasugai, T.Iida and A.Takahashi. Measurement of Formation Cross Cross Section of Short-Lived Nuclei by 14 MeV Neutrons, ibid.pp.361-363.
- [34] W. Mannhart, D. Schmidt and Xia Haihong, Measurement of the $^{59}\text{Co}(n,\alpha)^{56}\text{Mn}$, $^{59}\text{Co}(n,p)^{59}\text{Fe}$ and $^{59}\text{Co}(n,2n)^{58m+g}\text{Co}$ Cross Sections between 8 and 14 MeV, ibid., pp.285-287.
- [35] N.I.Molla, R.U.Miah, S.Basunia, S.M.Hossain and M.Rahman, Cross Sections of (n,p), (n, α) and (n,2n) Processes on Scandium, Vanadium, Cobalt, Copper and Zinc Isotopes in the Energy Range 13.57-14.71 MeV, ibid., pp.938-940.
- [36] V.N.Manokhin, A.B.Pashchenko, V.I.Plyaskin, V.M.Bychkov and V.G.Pronyaev, Activation Cross Sections Induced by Fast Neutrons, in Handbook on Nuclear Activation Data, Tech.Rep.Series No.273, IAEA, Vienna, 1987, pp.305-411.
- [37] F.Cserpak, S.Sudar, J.Csikai and S.M.Qaim, Excitation Functions and Isomeric Cross Section Ratios of the $^{63}\text{Cu}(n,\alpha)^{60m+g}\text{Co}$, $^{65}\text{Cu}(n,\alpha)^{62m+g}\text{Co}$ and $^{60}\text{Ni}(n,p)^{60m+g}\text{Co}$ Processes from 6-15 MeV, Phys.Rev., C49(1994)1525-1533.
- [38] I.G.Birn, B.Strohmaier, H.Freiesleben and S.M.Qaim, Isomeric Cross Section Ratios for the Formation of $^{75m+g}\text{Ge}$ in (n,p), (n, α) and (n,2n) Reactions from 6-15 MeV, Phys.Rev., C52(1995)2546-2551.
- [39] Y.Ikeda, E.T.Cheng, C.Konno and H.Maekawa, Measurement of Neutron Activation Cross Sections for the $^{99}\text{Tc}(n,p)^{99}\text{Mo}$, $^{99}\text{Tc}(n,\alpha)^{96}\text{Nb}$ and $^{99}\text{Tc}(n,n'\alpha)^{95}\text{Nb}$ and $^{99}\text{Tc}(n,n')^{99m}\text{Tc}$ Reactions at 13.5 and 14.8 MeV, Nucl.Sci.Eng.116(1994)28-34.
- [40] R. Klopris, R.Doczi, S.M.Qaim, S.Sudar and J.Csikai, Excitation Functions of some Neutron Threshold Reactions on ^{89}Y in the Energy Range of 7.8 to 14.7 MeV, Radiochimica Acta, 79(1997)3-9.
- [41] I.Birn and S.M.Qaim, Excitation Functions of Neutron Threshold Reactions on Some Isotopes of Germanium, Arsenic and Selenium in the 6.3-14.7 MeV Energy Range, Nucl.Sci.Eng.116(1994)125-137.
- [42] S.M.Qaim, Lack of Evidence for Shell Effects in the (n,p) and (n, α) Reaction Cross Sections at 14.7 MeV Neutron Energy, Z.f.Naturforschung, 25(1970)1977-1978.

- [43] Cai Dunjiu, Evaluation of Cross-Sections for Dosimetry Reactions, INDC(CPR)-024(October,1991), IAEA, Vienna.
- [44] Yuan Junqian, Kong Xiangzhong, Yang Jingkang and Wang Yongchang, Progress on 14 MeV Neutron Activation Cross Section Measurement at Lanzhou University, INDC(CPR)-032, CNDC-0014(1994)pp.5-8.
- [45] S. M. Qaim, St Spellerberg, F. Cserpak and J. Csikai, Radiochemical Measurement of Excitation Function of $^{63}\text{Cu}(\text{n},\text{p})^{63}\text{Ni}$ Reaction from 7.2 to 14.6 MeV, Radiochimica Acta, 73 pp. 111-117. (1996).
- [46] A. A. Filatenkov, S. V. Chuvaev, V. A. Jakovlev, A. V. Malyshenkov, S. K. Vasiljev, Systematic Measurement of Cross Sections at Neutron Energies of 13.4-14.9 MeV, NDST-97 May 19-24, 1997 Trieste.
- [47] S. V. Chuvaev, A. A. Filatenkov, V. A. Jakovlev, A. V. Malyshenkov, S. K. Vasiljev, M. Avrigeanu, V. Avrigeanu, Measurement and Analysis of the (n,p), (n,α), ($\text{n},2\text{n}$) and (n,np) Reaction Cross Sections of ^{59}Co and ^{58}Ni , NDST-97 May 19-24, 1997 Trieste.
- [48] Y. Kasugai, Y. Ikeda and Y. Uno, Activation Cross Section Measurement for La, Ce, Pr, Nd, Gd, Dy and Er isotopes by 14 MeV Neutrons, NDST-97 May 19-24, 1997 Trieste.
- [49] S. Amemiya, K. Ishibashi and T. Katoh, Neutron Activation Cross section of Molybdenum Isotopes at 14.8 MeV, Journal of Nuclear Science and Technology Vol. 19, No. 10, pp. 781-788, October 1982.
- [50] S. M. Qaim, Nuclear reaction Cross-Sections for 14.7 MeV Neutrons on ^{99}Tc , J. Inorg. Nucl. Chem., 1973, Vol. 35, pp. 3669-3675.
- [51] M. Herman, A. Marcinkowski and J. Bielewicz, and P. Oblozinsky, Cross Sections for the (n,p) and (n,n') Reactions on Pd, Cd and Os, Nuclear Physics A297 (1978) 335-346.
- [52] S. J. Hasan, A. Pavlik, G. Winkler, M. Uhl and M. Kaba, Precise measurement of cross sections for the reactions $^{59}\text{Co}(\text{n},2\text{n})^{58\text{m}+\text{g}}\text{Co}$ and $^{59}\text{Co}(\text{n},\text{p})^{59}\text{Fe}$ around 14 MeV, J. Phys. G: Nucl. Phys., 12 (1986) 397-410.
- [53] Y. Kasugai, H. Yamamoto, A. Takahashi, T. Iida and K. Kawade, Measurement of Activation Cross Sections on Ta, W for 14 MeV Neutrons, Jaeri-M 93-046.

- [54] I. Ribansky, Neutron Activation Cross Section Measurement at Bratislava, INDC (CSR)-008/GI INT (86)-2.
- [55] H. D. Luc, P. N. Ngoc, N. V. Do, L. B. Bach, Determination of some (n,p) (n,n'p) and (n, α) Reaction Cross sections Induced by 14.8 MeV Neutrons on Cr and Ti Isotopes, INDC (VN) -5/GI INT (86)-9.
- [56] K. T. Osman and F.I. Habbani, Measurement and Study of (n,p) Reaction Cross sections for Cr, Ti, Ni, Co, Zr and Mo Isotopes using 14.7 MeV Neutrons, INDC(SUD)-001 (1996).
- [57] Y. Junqian, K. Xiangzhong, Y. Jingkang, W. Huaiyi, cross Section Measurements for $^{199}\text{Hg}(\text{n},\text{p})^{199}\text{Au}$, $^{198}\text{Hg}(\text{n},\text{p})^{198}\text{Au}$, $^{196}\text{Hg}(\text{n},\text{p})^{196}\text{Au}$ and $^{196}\text{Hg}(\text{n},\text{d}+\text{np}+\text{pn})^{195}\text{Au}$ Reactions, Communication of Nuclear Data Progress No 13 (1995), INDC(CPRÖ-034/L), pp. 16-18.
- [58] A.B.Pashchenko, H.Wienke, J.Kopecky, J.Ch.Sublet and R.A.Forrest, Neutron Activation Cross Section Data Library for Fusion Applications, IAEA-NDS-173(March,1997).
- [59] Comparisson of Activation Cross Section Measurements and Experimental Techniques for Fusion Reactor Technology, (Ed. A.B.Pashchenko), INDC(NDS)-301(Julay,1994), INDC(NDS)-319(February,1995), INDC(NDS)-320(February,1995),IAEA,Vienna.
- [60] Activation Cross-Sections for the Generation of Long-Lived Radionuclides of Importance in Fusion Reactor Thechnology, (Ed. Wang Dahai), INDC(NDS)-232(January,1990), (Ed. A.B.Pashchenko), INDC(NDS)-286,288(November,1993), INDC(NDS)-340(November,1995), INDC(NDS)-342(February,1996), IAEA, Vienna.
- [61] Z.Bödy and J.Csikai, Data for 14 MeV Neutron Actvation Analysis, in Handbook of Nuclear Activation Data, Tech.Rep.Series, No.273, IAEA,Vienna(1987)pp.261-303.
- [62] Cs.M.Buczkó, J.Csikai, S.Sudár, A.Grallert, S.A.Jonah, B.W.Jimba, T.Chimoye, M.Wagner, Excitation Functions and Isotopic Cross Section of the $^{58}\text{Ni}(\text{n},\text{p})^{58}\text{Co}^{\text{m,g}}$ Reactions from 2 to 15 MeV, Phys.Rev.C52(1995)1940-1946.
- [63] J.Csikai, Nuclear Data Activities in Hungary, (Proceeding of the 1993 Symposium on Nuclear Data, JAERI-M, 94-019), INDC(JPN)-196, Tokai-mura (1994)pp.408-425.

- [64] J.Csikai and S.Nagy, Disintegration of ^{14}N by Fast Neutrons, Acta. Phys.Hungarica, 21(1966)303-310.
- [65] J.Csikai and S.Nagy, Some (n,p) Reaction Cross Sections for 14.7 MeV Neutrons, Nucl.Phys.91(1967)222-224.
- [66] Von M., Schmidt-Hönow and W.Herr, Die Aktivierungsschnitte der $^{16}\text{O}(\text{n},\text{p})^{16}\text{N}$ und $^{18}\text{O}(\text{n},\alpha)^{15}\text{C}$ Reactionen für 14 MeV (d,t) Neutronen, Radiochimica Acta, 17(1972) pp.142-146.
- [67] A.B.Lillie, The Disintegration of Oxigen and Nitrogen by 14.1 MeV Neutrons, Phys.Rev. 87(1952)716-722.
- [68] M.Bormann, S.Cierjacks, E.Fretwurst, K.J.Giesecke,H.Neuert and H.Pollehn, Untersuchungen über die Energieabhängigkeit von Kernreactionen mit Neutronen im Energiebereick zwischen 12 und 19 MeV, Z.f.Physik 174(1963)1-17.
- [69] H.Felber and H. Friedmann, Evaluation of the Cross Section $^{14}\text{N}(\text{n},\text{p})^{14}\text{C}$ for 14.7 MeV Neutrons by Measurement of the Activity of ^{14}C , Z.f.Physik, A276(1976)75-78.
- [70] M.E.Battat and F.L.Ribe, Formation of ^6He by 14 MeV Neutron Bombardment of Li and Be, Phys. Rev.89(1953)80-83.
- [71] P.Jessen, M. Bormann, F. Dreyer and H. Neuert, Experimental Excitation Functions for (n,p), (n,t), (n,α), (n,2n), (n,np) and (n,nα) Reactions, Nuclear Data, Academic Press, New York, London (1965) Vol. 1.,103-202.
- [72] K. Kawade, H. Yamamoto, T. Kobayashy, T. Katoh, T. Iida and A. Takahashi, Measurement of Formation Cross Sections of Short-Lived Nuclei by 14 MeV Neutrons, JAERI-M, 92-020, (1992), INDC(JPN)-154.
- [73] J. Csikai, J. Bacsó, A. Daróczy, Investigations of the Cross Section of Neutron Reactions in the Nucleus ^{103}Rh Nucl. Phys. 41(1963) 316-322.
- [74] R. Pepelnik, B. Anders, B.M. Bahal, Measurements of 14 MeV Neutron Activation Cross Sections, Radiation Effects, 92(1986) 211-214.
- [75] R. Wölflé, A. Mannan, S.M. Qaim and R. Videra, Excitation Functions of $^{93}\text{Nb}(\text{n},2\text{n})^{92m}\text{Nb}$, $^{93}\text{Nb}(\text{n},\alpha)^{90m,g}\text{Y}$, $^{139}\text{La}(\text{n},\alpha)^{136}\text{Cs}$ and $^{181}\text{Ta}(\text{n},\text{p})^{181}\text{Hf}$ Reactions in the Energy Range of 12.5-19.6 MeV, Applied Rad. Isot.39(1988) 407-412.
- [76] J. Csikai, Activation Cross Sections for the Generation of Long-Lived Radionuclides of Importance in Fusion Reactor Technology, [Ed. A.B. Paschenko] INDC(NDS-342) 29-35, IAEA, Vienna, (February, 1996).

- [77] Xiangzhong Kong, Yongchang Wang, Jingkang Yang and Junqian Yuan, Cross Sections for (n,p), (n, α) and (n,2n) Reactions in Rare-Earth Isotopes at 14.7 MeV, *Radiochimica Acta* (in press).
- [78] Zhao Wenrong, Yu Weixiang and Lu Hanlin, Cross Section Measurements for Reactions at Neutron Energies of 9.5, 9.9 and 14.7 MeV, CIAE Annual Report (1994) pp.23-24.
- [79] Y.Satoh, T. Matsumoto, Y. Kasugai, H. Yamamoto, T. Iida, A. Takahashi and K. Kawade, Measurement of Formation Cross Sections Producing Short-Lived Nuclei by 14 MeV Neutrons, -Na, Si, Te, Ba, Ce, Sm, W, Os-, Proceeding of the 1994 Symposium on Nuklear Data, JAERI Conf-95-008, pp.189.
- [80] K. Sailer, S. Daróczy, P. Raics and S. Nagy, The Cross Sections of (n,2n), (n,p) and (n, α) Reactions for 14 MeV Neutrons on Isotopes of Cr. and Zr, *Neytronnaya Fizika* 1(1977)276, Moskwa.
- [81] V. N. Levkovskii, Empirical behaviour of the (n,p) cross section for 14-15 MeV Neutrons, *Zh. Eksp. Teor. Fiz.* 45,(1963) 305-311.
- [82] V.N. Levkovskii, Cross Sections of (n,p) and (n, α) Reactions at the Neutron Energy of 14-15 MeV, *Jadernaja Physica*, 18(1973) 705-709.
- [83] G. Eder, G. Wnkler and P. Hille, *Z. Phys.* 253, (1972), 355.
- [84] V. M. Bychkov, V. N. Manokin, A. B. Paschenko and V. I. Plyaskin, (1980) INDC (CCP) 146-LJ, 1.
- [85] S Ait-Tahar, The systemstics of (n,p)cross sections for 14 MeV neutrons, *Nucl. Phys.* 13.(1987) pp.121-125.
- [86] S. Selvi and H. H. Erbil, Empirical formulae for 14.5 MeV (n,p) cross sections, INDC(TUR)-002/L.
- [87] A. I. Dityuk, A Yu. Konobeyev, V. P. Lunev, Yu. N. Shubin, New Approach for the Creation of Neutron Threshold Reaction Systematics, NDST-97, May 19-24, 1997 Trieste.
- [88] N. Yamamuro, A Nuclear Cross Section Calculation System with Simplified Input-Format VERSION-II (SINCROS-II), JAERI-M 90-006, Japan Atomic Energy Research Institute (1990).
- [89] N. Yamamuro, Cross-Section Calculations for Neutron-Induced Reactions up to 50 MeV, *Nucl. Sci. and Eng.*,122(1996) 374-383.

- [90] N. Yamamuro, Calculation of Activation Cross Sections for Molybdenum Isotopes, Nucl. Sci. and Eng., 109(1991) 128-141.
- [91] N. Yamamuro, Activaton Cross-Section Calculations on the Production of Long-Lived Radionuclides, Nucl. Sci. and Eng., 118(1994) 249-259.

Table 1. Some recommended (n,p) cross sections at 14.7 ± 0.2 MeV neutron energy

Target Element	Z	A	$\sigma \pm \Delta\sigma$ (mb)
B	5	11	4.2 ± 0.2
N	7	14	45.2 ± 6
O	8	18	2.3 ± 0.5
F	9	19	18.6 ± 0.88
Na	11	23	47.0 ± 2
Mg	12	24 25 26	183.6 ± 9 88.04 ± 13 39 ± 11
Al	13	27	70 ± 2
Si	14	28 29 30	238.7 ± 30 135.33 ± 15 70.04 ± 20
P	15	31	91.85 ± 3.48
S	16	32 33 34	237 ± 25 134 ± 22 78 ± 8
Cl	17	35 37	110 ± 10 25.4 ± 3
K	19	39 41	314 ± 14 51.36 ± 3
Ca	20	40 42 43 44	470.01 ± 50 178.25 ± 12 99.83 ± 13 43.27 ± 6
Sc	21	45	57 ± 8
Ti	22	46 47 48 50	251 ± 13 136 ± 9 57.2 ± 2.7 11.9 ± 6
V	23	51	33.3 ± 1.7
Cr	24	50 52 53 54	294 ± 30 80 ± 4 42 ± 2.1 17.4 ± 1
Mn	25	55	45 ± 10
Fe	26	54 56 57	350 ± 15 115 ± 6 59 ± 4

Table 1. Cont.

Co	27	59	51±1
Ni	28	58	366±19
		60	148±8
		61	64±4
		62	37±4
Cu	29	63	70±13
		65	20.93±4
Zn	30	64	185±10
		66	73.3±8
		67	37.9±6
		68	15.2±2.5
Ga	31	69	64±5
		71	20.5±1.5
Ge	32	70	74±8
		72	31.6±1.4
		73	19.4±1.1
		74	10.1±0.5
		76	2.8±0.4
As	33	75	18.7±2.1
Se	34	74	112±7
		76	49±3
		78	19±1.1
Br	35	81	21±5
Kr	36	80	45±6
		82	23±5
		84	11±2
		86	5±1.5
Sr	38	84	95±8
		86	44±4
		88	17.4±2
Y	39	89	22.2±4.2
Zr	40	90	37±5
		91	29±4
		92	20.23±2.5
		94	7.5±1.1
Mo	42	95	41.3±2
		96	21.3±1.1
		97	14.6±0.8
Ru	44	96	146±8
		101	21.2±1.2
		104	6±0.7
Rh	45	103	17±3
Pd	46	102	93.6±15
		104	58±15
		105	38±2.9

Table 1. Cont.

Pd	46	106 108	22.5 ± 6 4 ± 1
Cd	48	106	130 ± 24
		111	23.25 ± 2.1
		112	16.1 ± 3
		114	8.5 ± 1.3
		116	2.96 ± 0.3
In	49	115	13.26 ± 2.95
Sn	50	116	14.6 ± 2
		117	11.7 ± 0.6
		119	7.1 ± 0.8
		120	4.5 ± 0.5
Te	52	122	10.5 ± 1.5
		126	4.7 ± 0.2
		128	2.9 ± 0.1
		130	1.7 ± 0.1
I	53	127	11.7 ± 2
Xe	54	128	27 ± 4
		131	6.1 ± 0.6
Cs	55	133	10.5 ± 2
Ba	56	132	15.3 ± 3.5
		136	4.8 ± 0.7
		138	2.18 ± 0.15
La	57	139	4.5 ± 1.1
Ce	58	140	7.05 ± 0.7
		142	4.8 ± 0.8
Pr	59	141	11.5 ± 0.9
Nd	60	142	13.7 ± 1.1
		143	11.5 ± 2.3
		144	9.8 ± 1.5
		145	7.25 ± 1.6
		146	4.5 ± 0.7
Sm	62	144	19 ± 4
		148	9.8 ± 0.8
		150	7 ± 0.6
Eu	63	153	4.2 ± 0.4
Gd	64	157	5.4 ± 1.1
		158	3.2 ± 1.2
Tb	65	159	5.1 ± 0.4
Dy	66	158	10.6 ± 1.2
		160	7 ± 1.5
		162	4.3 ± 1
		164	2.55 ± 0.5
Er	68	166	4.5 ± 0.7
		167	3.4 ± 0.3

Table 1. Cont.

Er	68	168	2.8 ± 0.4
Yb	70	174	3 ± 0.2
Lu	71	175	4 ± 0.7
Ta	73	181	2.94 ± 0.18
W	74	182	6.5 ± 0.5
		184	2.61 ± 0.12
		186	1.25 ± 0.25
Re	75	187	3.73 ± 0.28
Os	76	184	9.30 ± 2
		188	4 ± 0.8
		190	2.2 ± 0.5
		192	1 ± 0.03
Au	79	179	2 ± 0.5
Hg	80	196	9.3 ± 1
		198	4.5 ± 0.8
		199	2.5 ± 0.5
Tl	81	203	4.2 ± 0.8
		205	1.9 ± 0.2
Bi	83	209	0.8 ± 0.3

Table 2. Deduced (n,p) cross sections using Eq. 3 at 14.7 ± 0.2 MeV

Target Element	Z	A	σ (mb)
S	16	36	28.6
Ca	20	46	9.2
		48	1.5
Ti	22	49	23.1
Fe	26	58	28.4
Ni	28	64	3.75
Zn	30	70	2.15
Se		77	30.9
	34	80	6.7
		82	2.16
Kr	36	78	81.4
		83	16
Sr	38	87	28.4
Zr	40	96	1.93
Mo		92	186.7
	42	94	69.4
		98	7
		100	1.96
Ru		98	68.1
	44	99	46.3
		100	31.3
		102	14.1
Pd	46	110	0.82
Cd		108	66.7
	48	110	33.3
		113	11.3
Sn		112	41.6
		114	25
		118	8.7
		122	2.9
		124	1.68

Target Element	Z	A	σ (mb)
Te		120	14.3
	52	123	8.8
		124	7.2
		125	5.9
Ba	56	130	24.8
Ba	56	134	8.7
		135	6.4
Nd	56	137	3.3
	60	148	3.3
Sm		150	1.9
		147	11.6
	62	149	8.3
		152	5
Dy		154	3.5
		156	15.2
	66	161	5.5
Er		163	3.3
		162	11.4
	68	164	7.2
W		170	1.66
	74	180	15.2
Os		183	4.2
		186	5.96
	76	187	4.7
Hg		189	2.9
		200	1.85
	80	201	1.22
		202	0.8
		204	0.34

Table 3. Total (n,p) cross sections for elements at (14.7±0.2) MeV

Element	Z	σ_{tot} (mb)
N	7	45.2±6
O	8	2.3±0.5
F	9	18.6±0.9
Na	11	47.0±2
Mg	12	158.0±10
Al	13	70.0±2
Si	14	231.4±30
P	15	91.9±3.5
S	16	229.5±25
Cl	17	89.5±10
K	19	296.3±14
Ca	20	457.8±50
Sc	21	57.0±8
Ti	22	74.9±12
V	23	33.3±1.7
Cr	24	84.2±18
Mn	25	45.0±10
Fe	26	127.3±15
Co	27	51.0±1
Ni	28	290.1±19
Cu	29	54.9±13
Zn	30	114.7±10
Ga	31	46.6±5
Ge	32	29.8±8
As	33	18.7±2.1
Se	34	16.0±4
Kr	36	13.0±3

Element	Z	σ_{tot} (mb)
Sr	38	21.2±5
Y	39	22.2±4.2
Zr	40	27.1±5
Mo	42	47.5±2
Ru	44	28.3±4
Rh	45	17.0±3
Pd	46	23.2±4
Cd	48	17.2±5
Sn	50	8.1±2
Te	52	3.5±1
I	53	11.7±2
Cs	55	10.5±2
Ba	56	3.0±0.7
La	57	4.5±1.1
Pr	59	11.5±0.9
Nd	60	9.1±1.5
Sm	62	7.2±1
Tb	65	5.1±0.4
Dy	66	3.9±1
Er	68	3.4±0.7
Ta	73	2.9±0.2
W	74	3.5±0.5
Os	76	2.2±0.6
Au	79	2.0±0.5
Hg	80	1.7±0.5
Tl	81	2.6±0.3
Bi	83	0.8±0.3

Table 4. Comparison of (n,p) systematics at (14.7±0.2) MeV.

Author	Formula, σ (mb)	Mass region	n*	F/n
Levkovskii	$\sigma = \begin{cases} 55.3(A^{1/3} + 1)^2 \exp\left(-34.8 \frac{N - Z + 1}{A + 1}\right) \\ 49.4(A^{1/3} + 1)^2 \exp\left(-35.1 \frac{N - Z}{A}\right) \end{cases}$	19≤A≤40	17	0.468
		40≤A≤188	175	0.399
Eder et al.	$\sigma = \exp\left(1.31 + 0.806A^{1/2} - 10.3 \frac{N - Z}{A^{2/3}}\right)$	19≤A≤188	191	0.29
Bychkov et al.	$\sigma = 46.6(A^{1/3} + 1)^2 \exp\left\{\sqrt{\frac{A}{140}}\left(-53.3 \frac{N - Z + 1}{A} + 0.622 \frac{Z - 1}{A^{1/3}} - 3.20\right)\right\}$	40≤A≤188	175	0.307
Forrest	$\sigma = 11.23(A^{1/3} + 1)^2 \exp\left(-32.73 \frac{N - Z}{A} - 46.57\left(\frac{N - Z}{A}\right)^2 + 0.218A^{1/2}\right)$	40≤A≤188	175	0.254
Kumabe and Fukuda	$\sigma = \begin{cases} 27.9A \exp\left(-39.1 \frac{N - Z}{A}\right) \\ 0.58A^2 \exp\left(-42.3 \frac{N - Z}{A}\right) \\ 0.94A^2 \exp\left(-47.8 \frac{N - Z}{A}\right) \end{cases}$	40≤A≤62	25	0.200
		63≤A≤89	28	0.336
		90≤A≤188	113	0.289
Ait-Tahar	$\sigma = 140.2(A^{1/3} + 1)^2 \exp\left(-39.1 \frac{N - Z + 1}{A}\right)$	40≤A≤188	175	0.309

Table 4. Cont.

Author	Formula, σ (mb)	Mass region	n*	F/n
Kasugai et al.	$\sigma = 1830(N - Z + 1) \exp\left(-50.7 \frac{N - Z + 1}{A}\right)$	$19 \leq A \leq 188$	191	0.280
Korovin et al.	$\sigma = \pi r_0^2 \left(A^{1/3} + 1\right)^2$ $\times \left\{ A^{1.1128} \left(1.1242 \left(\frac{N - Z + 1}{A}\right)^2 - 0.73212 \left(\frac{N - Z + 1}{A}\right) + 0.11707 \right)^3 \right.$ $\left. + 0.4936 \exp\left(-194.69 \left(\frac{N - Z + 1}{A}\right)^2 - 5.3778 \left(\frac{N - Z + 1}{A}\right)\right) \right\}$	$11 \leq A \leq 209$	207	0.331
Present	$23.659 \left(A^{1/3} + 1\right)^2 \exp\left(-23.041 \left(\left(\frac{N - Z}{A}\right) + \left(\frac{N - Z}{A}\right)^2\right)\right)$	$11 \leq A \leq 209$	207	0.351

*Number of data points

Table 5. Estimated (n,p) cross sections for long-lived target and residual nuclei at 14.7 ± 0.2 MeV.
 $T_{1/2}$ gives target $T_{1/2}/\text{residual } T_{1/2}$ in years (if not stated otherwise).

Reaction	$T_{1/2}$	σ (mb)
$^{40}\text{Ca(n,p)}^{40}\text{K}$	$\infty/1.28 \cdot 10^9$	470 ± 50
$^{50}\text{Cr(n,p)}^{50}\text{V}$	$\infty/1.4 \cdot 10^{17}$	294 ± 50
$^{60}\text{Ni(n,p)}^{60}\text{Co}$	$\infty/5.27$	148 ± 8
$^{63}\text{Cu(n,p)}^{63}\text{Ni}$	$\infty/100.1$	70 ± 13
$^{92}\text{Mo(n,p)}^{92}\text{Nb}$	$\infty/3.47 \cdot 10^7$	186.7
$^{94}\text{Mo(n,p)}^{94}\text{Nb}$	$\infty/2.03 \cdot 10^4$	69.4
$^{108}\text{Cd(n,p)}^{108}\text{Ag}$	$\infty/2.37\text{m} + 418$	66.7
$^{134}\text{Ba(n,p)}^{134}\text{Cs}$	$\infty/2.062$	8.7
$^{135}\text{Ba(n,p)}^{135}\text{Cs}$	$\infty/2.3 \cdot 10^6 + 53\text{m}$	6.4
$^{137}\text{Ba(n,p)}^{137}\text{Cs}$	$\infty/30.07$	3.3
$^{158}\text{Dy(n,p)}^{158}\text{Tb}$	$\infty/180$	10.6 ± 1.2
$^{180}\text{W(n,p)}^{180}\text{Ta}$	$\infty/8.15\text{h} + >1.2 \cdot 10^{15}$	15.2
$^{32}\text{Si(n,p)}^{32}\text{Al}$	$172/33\text{ms}$	16
$^{41}\text{Ca(n,p)}^{41}\text{K}$	$1.03 \cdot 10^5/\infty$	263
$^{55}\text{Fe(n,p)}^{55}\text{Mn}$	$2.73/\infty$	145
$^{59}\text{Ni(n,p)}^{59}\text{Co}$	$7.6 \cdot 10^4/\infty$	165
$^{60}\text{Fe(n,p)}^{60}\text{Mn}$	$1.5 \cdot 10^6/51\text{s} + 1.77\text{s}$	17
$^{63}\text{Ni(n,p)}^{63}\text{Co}$	$100.1/27.4\text{s}$	34
$^{79}\text{Se(n,p)}^{79}\text{As}$	$\leq 6.5 \cdot 10^4/9.01\text{m}$	17
$^{81}\text{Kr(n,p)}^{81}\text{Br}$	$2.29 \cdot 10^5/\infty$	39
$^{85}\text{Kr(n,p)}^{85}\text{Br}$	$10.756/2.90\text{m}$	12
$^{90}\text{Sr(n,p)}^{90}\text{Rb}$	$28.78/158\text{s} + 258\text{s}$	11
$^{93}\text{Zr(n,p)}^{93}\text{Y}$	$1.53 \cdot 10^6/10.18\text{h} + 0.82\text{s}$	18
$^{93}\text{Mo(n,p)}^{93}\text{Nb}$	$4.0 \cdot 10^3/\infty + 16.13\text{y}$	63
$^{107}\text{Pd(n,p)}^{107}\text{Rh}$	$6.5 \cdot 10^6/21.7\text{m}$	20
$^{126}\text{Sn(n,p)}^{126}\text{In}$	$\approx 1 \cdot 10^5/1.6\text{s}$	2.7
$^{133}\text{Ba(n,p)}^{133}\text{Cs}$	$10.52/\infty$	13
$^{146}\text{Sm(n,p)}^{146}\text{Pm}$	$10.3 \cdot 10^8/5.53$	17
$^{151}\text{Sm(n,p)}^{151}\text{Pm}$	$90/28.40\text{h}$	7.4
$^{154}\text{Dy(n,p)}^{154}\text{Tb}$	$3.0 \cdot 10^6/21.5\text{h} + 9.4\text{h} + 22.7\text{h}$	22
$^{194}\text{Hg(n,p)}^{194}\text{Au}$	$520/38.02\text{h} + 600\text{ms} + 420\text{ms}$	9.5

Fig 1. Isotopic dependence of $\sigma_{n,p}$ data





Fig 2. The value of $\sigma_{n,p} / (A^{1/3} + 1)^2$ as a function of $(N-Z)/A$







Fig. 3a The distribution of the deviations $((\sigma_{\text{exp}} - \sigma_{\text{calc}})/\sigma_{\text{calc}})$ for (n,p) reactions.




Fig. 3b The distribution of the deviations $((\sigma_{\text{exp}} - \sigma_{\text{calc}})/\sigma_{\text{calc}})$ for (n,p) reactions.




Fig. 3c The distribution of the deviations $((\sigma_{\text{exp}} - \sigma_{\text{calc}})/\sigma_{\text{calc}})$ for (n,p) reactions.

Nuclear Data Section
International Atomic Energy Agency
P.O. Box 100
A-1400 Vienna
Austria

e-mail: services@iaeand.iaea.or.at
fax: (43-1)20607
cable: INATOM VIENNA
telex: 1-12645 atom a
telephone: (43-1)2060-21710

online: TELNET or FTP: iaeand.iaea.or.at
username: IAEANDS for interactive Nuclear Data Information System
username: ANONYMOUS for FTP file transfer
For users with Web-browsers: <http://www-nds.iaea.or.at>
