

# Benchmark of Spallation Models



## Benchmark objectives

Sylvie LERAY

CEA/Saclay, IRFU/SPhN

# Spallation reaction applications

- Spallation neutron sources
- ADS
- RIB production
- Detection setup simulation
- Radioprotection near accelerators
- Hadrontherapy
- Astrophysics
- Cosmic rays in space



MYRRHA

# Necessity for a benchmark

- Large amount of high quality data collected
  - ↳ Neutron and light charged particle production, isotopic residue distributions, excitation functions
- ≈ HINDAS FP5 and EUROTRANS/NUDATRA FP6 projects
- Improvement of nuclear models
  - ↳ INCL4/ABLA, FLUKA, CEM....
  - ↳ new approaches: QMD models, BUU, VUU....
- Implementation into high-energy transport codes
  - ↳ examples: INCL4/ABLA and CEM into MCNPX, BIC and INCL4/ABLA in GEANT4, JQMD in PHITS...

# Joint ICTP-IAEA Advanced Workshop on Model Codes for Spallation Reactions (Trieste, February 2008)

## Goals

- To bring together experts on spallation models
- To understand in depth, the physics of INC, QMD models and de-excitation models
- To define an agreed set of experimental data to be used for the benchmarking of the models
- To define the specifications of the benchmark



The Abdus Salam  
International Centre for Theoretical Physics

### Joint ICTP-IAEA Advanced Workshop on Model Codes for Spallation Reactions

4 - 8 February 2008  
(Miramare, Trieste, Italy)

The International Atomic Energy Agency (IAEA) together with the Abdus Salam International Centre for Theoretical Physics (ICTP), will organize an *Advanced Workshop on Model Codes for Spallation Reactions* which will be held at the ICTP, Trieste, Italy, from 4 - 8 February 2008.

Spallation reactions play an important role in a wide domain of applications ranging from intense neutron sources for condensed matter and material studies, accelerator-driven sub-critical reactors for the transmutation of nuclear waste and rare isotope production to astrophysics, simulation of detector set-ups in nuclear and particle physics experiments, and radiation protection near accelerators or in space. The simulation tools in these domains are highly complex, typically codes in several parts. In particular, predictions and characterizations of all reaction products are taken from existing experimental library data or, when experimental data are missing, calculated using nuclear model codes as event generators. These are generally Monte-Carlo implementations of Intra-Nuclear Cascade (INC) models or Quantum Molecular Dynamics (QMD) models followed by de-excitation (principally evaporation/fission) models. It is of great importance to validate abilities of the various codes to predict reliably, with a known uncertainty, the different quantities relevant for applications.

**PURPOSE:** This Workshop will facilitate experts and competent practitioners to better understand the physical basis, approximations, strengths and weaknesses of the currently used spallation codes. Presentation of relevant basic experimental data with emphasis on accuracies, detector efficiencies, filters and thresholds will create basis for code validation and inter-comparison. Specifically the workshop will help:

- \* To understand in depth, the physics of INC, QMD models and de-excitation models to point out the reasons of their respective successes or deficiencies;
- \* To define an agreed set of experimental data to be used in validation and inter-comparison of the models;
- \* To promote the exchange of information among researchers in the field;
- \* To identify areas of international cooperation in the field.

The agreed set of experimental data will be proposed as an international benchmark and reviewed by experts in a follow-up activity.

**PARTICIPATION:** Experts, young scientists, and Ph.D. students from all countries which are members of the United Nations, UNESCO or IAEA may attend the advanced workshop. As it will be conducted in English, participants should have an adequate working knowledge of this language. Also, it is intended that the conference will attract research workers from developing countries, through a programme of training activities within a framework of international cooperation, a limited number of young scientists, Ph.D. students, and post-doctoral scientists from developed countries are also welcome to attend this Workshop.

As a rule, travel and subsistence expenses of the participants should be borne by the home institution. Every effort should be made by candidates to secure support for their fare (or at least half-fare). However, limited funds are available for some participants, who are nationals of, and working in, a developing country. Preference will be given to qualified candidates not more than 45 years old. Such support is available only for those who attend the entire activity. There is no registration fee.


**Requests for Participation:** The 'Request for Participation' form is obtainable via Web server: <http://agenda.ictp.it/smri.php?1930>. It should be completed, signed and returned by using only one of the following ways:

If sending an application by e-mail to: [smr1930@ictp.it](mailto:smr1930@ictp.it) please save and send file attachments in either PDF (preferably) or RTF zipped or DOC format.

If sending an application form by regular mail or courier it should be posted to:  
Joint ICTP-IAEA Advanced Workshop on Model Codes for Spallation Reactions  
(smr1930 c/o Ms. Patricia Wardell)  
The Abdus Salam International Centre for Theoretical Physics  
Strada Costiera 11  
34014 Trieste, Italy (recent photograph & signature of the candidate are compulsory)

**ACTIVITY SECRETARIAT:** Telephone: +39-040-2240576      Telefax: +39-040-2240585  
E-mail: [smr1930@ictp.it](mailto:smr1930@ictp.it)      ICTP Home Page: <http://www.ictp.it/>

Trieste, September 2007



#### CO-SPONSORS:

CEA (Saclay, France)  
FZJ (Juelich, Germany)

#### DIRECTORS:

G. MANK  
(IAEA, Vienna, Austria)

D. FILGES  
(FZJ, Juelich, Germany)

S. LERAY  
(CEA, Saclay, France)

Y. YARIV  
(Soreq NRC, Yavne, Israel)

#### LOCAL ORGANIZER:

C. TUNIZ  
(ICTP, Trieste, Italy)

#### TOPICS:

INC Models

QMD and BUU Models

De-Excitation Models

Spallation reaction data for validation of models

Model inter-comparison methods

#### DEADLINE for requesting participation

**2 NOVEMBER 2007**

# Specifications of the Benchmark

- Participants should treat the complete reaction
  - complete reaction description (INC/QMD + De-ex)
- Participants should calculate the whole mandatory set of experimental data
  - + additional set if they have enough time
- Participants should give a comprehensive description of ingredients and parameters
  - list of the main ingredients and parameters
  - additional information requested ( $E^*$ ,  $A_R$ , ...)
- Calculations with one model should be done with the same set (default) parameters
  - predictive power
- Participants should give the source code

# Specifications

- **Domain: N + A, 20 MeV to 3 GeV, A >11**  
**why 20 MeV?**
  - 20-150 MeV libraries not available for all isotopes
  - for residue production below 150 MeV
  - to calculate correlations between particles
- **Set of data to be used for the benchmark**
  - All reaction channels
  - Mandatory data set : restriction to a limited number of systems (mainly Fe and Pb), a few energies covering the full range



# Neutron production

## Double-differential Cross Sections

| Beam | Target | Energy [MeV] | Angles [degrees] | Laboratory            | Reference   | Data  |
|------|--------|--------------|------------------|-----------------------|---|---|
| n    | Fe     | 65           | 9.5 - 28         | Univ. California, USA | E. L. Hjort et al., Phys. Rev. C 53 (1996) 237 -            | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| p    | Fe     | 800          | 0 - 160          | Saturne, France       | S. Leray et al., Phys. Rev. C 65 (2002) 044621 -            | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
|      |        |              | 30 - 150         | LANL, USA             | W.B. Amian et al., Nucl. Sci. Eng. 112 (1992) 78 -          | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| p    | Fe     | 1200         | 0 - 160          | Saturne, France       | S. Leray et al., Phys. Rev. C 65 (2002) 044621 -            | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| p    | Fe     | 1600         | 0 - 160          | Saturne, France       | S. Leray et al., Phys. Rev. C 65 (2002) 044621 -            | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| p    | Fe     | 3000         | 15 - 150         | KEK, Japan            | K. Ishibashi et al., J. of Nucl. Sci. Tech. 34 (1997) 529 - | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| p    | Pb     | 63           | 24 - 140         | Louvain, Belgium      | A. Guertin et al., Eur. Phys. J. A23 (2005) 49 -            | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| p    | Pb     | 256          | 8 - 150          | LANL, USA             | M. Meier et al., Nucl. Sci. Eng. 110 (1993) 289             | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| p    | Pb     | 800          | 0 - 160          | Saturne, France       | S. Leray et al., Phys. Rev. C 65 (2002) 044621 -            | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
|      |        |              | 30 - 150         | LANL, USA             | W.B. Amian et al., Nucl. Sci. Eng. 112 (1992) 78 -          | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| p    | Pb     | 1200         | 0 - 160          | Saturne, France       | S. Leray et al., Phys. Rev. C 65 (2002) 044621 -            | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| p    | Pb     | 1600         | 0 - 160          | Saturne, France       | S. Leray et al., Phys. Rev. C 65 (2002) 044621 -            | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| p    | Pb     | 3000         | 15 - 150         | KEK, Japan            | K. Ishibashi et al., J. of Nucl. Sci. Tech. 34 (1997) 529 - | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |

## Multiplicity Distributions


| Beam | Target | Energy [MeV]    | Laboratory      | Reference  | Data  |
|------|--------|-----------------|-----------------|--|---|
| p    | Fe     | 800, 1200, 1600 | Saturne, France | S. Leray et al., Phys. Rev. C 65 (2002) 044621 -           | <a href="#">data</a>                          |
| p    | Fe     | 1200            | COSY, Germany   | C.-M. Herbach et al., Jülich annual report (2001) -        | <a href="#">data</a>   <a href="#">figure</a> |
| p    | Pb     | 800, 1200, 1600 | Saturne, France | S. Leray et al., Phys. Rev. C 65 (2002) 044621 -           | <a href="#">data</a>                          |
| p    | Pb     | 1200            | COSY, Germany   | A. Letourneau et al., Nucl. Inst. Meth. B 170 (2000) 299 - | <a href="#">data</a>   <a href="#">figure</a> |



## Light charged particle production


### Double-differential Cross Section

| Beam | Target | Energy [MeV] | Emitted particles                 | Laboratory                | Reference  | Data   |
|------|--------|--------------|-----------------------------------|---------------------------|--|--|
| n    | Bi     | 542          | p, d, t                           | PSI, Switzerland          | J. Franz et al., Nucl. Phys. A 510 (1990) 774 -            | <a href="#">data</a>   <a href="#">figure</a>   EXFOR (p - d - t)                              |
| p    | Al     | 160          | $\alpha$                          | iTHEMBA, South Africa     | A. Cowley et al., Phys. Rev. C 54 (1996) 778 -             | <a href="#">data</a>   <a href="#">figure</a>   EXFOR  |
| p    | Fe     | 62           | p, d, t, $^3\text{He}$ , $\alpha$ | LANL, USA                 | F.E. Bertrand and R.W. Pelle, Phys. Rev. C 8 (1973) 1045 - | <a href="#">data</a>   <a href="#">figures</a>   EXFOR (p - d - t - $^3\text{He}$ - $\alpha$ ) |
| p    | Ni     | 175          | p, d, t, $^3\text{He}$ , $\alpha$ | COSY, Germany             | F. Goldenbaum et al. (unpublished)                         | <a href="#">data</a>   <a href="#">figures</a>   |
|      |        |              | p                                 |                           | S.V. Förtsch et al., Phys. Rev. C 43 (1991) 691 -          | <a href="#">data</a>   <a href="#">figure</a>   EXFOR  |
| p    | Ta     | 1200         | p, d, t, $^3\text{He}$ , $\alpha$ | COSY, Germany             | C.-M. Herbach et al., Nucl. Phys. A 765 (2006) 426 -       | <a href="#">data</a>   <a href="#">figure</a>  |
| p    | Au     | 160          | $\alpha$                          | iTHEMBA, South Africa     | A. Cowley et al., Phys. Rev. C 54 (1996) 778 -             | <a href="#">data</a>   <a href="#">figure</a>   EXFOR  |
| p    | Au     | 1200         | p, d, t, $^3\text{He}$ , $\alpha$ | COSY, Germany             | A. Budzanowski et al. (to be published)                    | <a href="#">data</a>   <a href="#">figures</a>   |
| p    | Au     | 2500         | p, d, t, $^3\text{He}$ , $\alpha$ | COSY, Germany             | A. Letourneau et al., Nucl. Phys. A 712 (2002) 133 -       | <a href="#">data</a>   <a href="#">figures</a>   |
|      |        |              |                                   |                           | A. Bubak et al., Phys. Rev. C 76 (2007) 014618 -           | <a href="#">data</a>   <a href="#">figures</a>   |
| p    | Pb     | 63           | p, d, t, $^3\text{He}$ , $\alpha$ | Louvain-la-Neuve, Belgium | A. Guertin et al., Eur. Phys. J. A 23 (2005) 49 -          | <a href="#">data</a>   <a href="#">figure</a>   EXFOR (p - d - t - $^3\text{He}$ - $\alpha$ )  |
| p    | Pb     | 800          | p                                 | LANL, USA                 | R. Chrien et al., Phys. Rev. C 21 (1980) 1014 -            | <a href="#">data</a>   <a href="#">figure</a>   EXFOR  |
|      |        |              |                                   |                           | J.A. McGill et al., Phys. Rev. C 29 (1984) 204 -           | <a href="#">data</a>   <a href="#">figure</a>   EXFOR  |
| p    | Bi     | 62           | p, d, t, $^3\text{He}$ , $\alpha$ | LANL, USA                 | F.E. Bertrand and R.W. Pelle, Phys. Rev. C 8 (1973) 1045 - | <a href="#">data</a>   <a href="#">figures</a>   EXFOR (p - d - t - $^3\text{He}$ - $\alpha$ ) |




### Pion production

| Beam | Target | Energy [MeV] | Emitted particles | Laboratory | Reference   | Data  |
|------|--------|--------------|-------------------|------------|---|---|
| p    | C      | 730          | $\pi^+$ , $\pi^-$ | LANL, USA  | D. R. F. Cochran et al., Phys. Rev. D 6 (1972) 3085 - | <a href="#">data</a>   <a href="#">figure</a> |
| p    | Al     | 730          | $\pi^+$ , $\pi^-$ | LANL, USA  | D. R. F. Cochran et al., Phys. Rev. D 6 (1972) 3085 - | <a href="#">data</a>   <a href="#">figure</a> |
| p    | Cu     | 730          | $\pi^+$ , $\pi^-$ | LANL, USA  | D. R. F. Cochran et al., Phys. Rev. D 6 (1972) 3085 - | <a href="#">data</a>   <a href="#">figure</a> |
| p    | Pb     | 730          | $\pi^+$ , $\pi^-$ | LANL, USA  | D. R. F. Cochran et al., Phys. Rev. D 6 (1972) 3085 - | <a href="#">data</a>   <a href="#">figure</a> |
| p    | Al     | 2205         | $\pi^-$           | KEK, Japan | H. En'yo et al., Phys. Lett. 159B (1985) 1 -          | <a href="#">data</a>   <a href="#">figure</a> |




## Production cross-sections from threshold to 3 GeV (excitation functions)

| Beam | Target | Energy [MeV] | Laboratory  | Reference   | Data   |
|------|--------|--------------|---|---|--|
| p    | Fe     | 20 to 3000   | Hannover University, Germany<br>University of Bern, Switzerland<br>ITEP, Russian Federation | R. Michel et al., Nucl. Sci. Tech., Supplement 2 (2002) 242 -<br>K. Ammon, I. Leya et al., Nucl. Instr. and Meth. B 266 (2008) 2 -<br>R. Michel et al., Nucl. Instr. and Meth. B 103 (1995) 183 -;<br>Th. Schiekel, R. Michel et al., Nucl. Instr. and Meth. B 114 (1996) 91 -;<br>R. Michel et al., Nucl. Instr. and Meth. B 129 (1997) 153 -<br>Titarenko (soon, october 2008?) | <a href="#">data</a>   <a href="#">figure</a><br><a href="#">data</a>   <a href="#">figure</a><br><br><a href="#">data</a>   <a href="#">figure</a><br><br><a href="#">data</a>   <a href="#">figure</a> |
| p    | Pb     | 20 to 3000   | ITEP, Russian Federation<br>ETH Zürich, Switzerland<br>Hannover University, Germany         | Y. E. Titarenko et al., Nucl. Instr. and Meth. A562 (2006) 801 -<br>M. Gloris et al., Nucl. Instr. and Meth. A463 (2001) 593 -<br>I. Leya et al., Nucl. Instr. and Meth. B229 (2005) 1 -  | <a href="#">data</a>   <a href="#">figure</a><br><a href="#">data</a>   <a href="#">figure</a><br><a href="#">data</a>   <a href="#">figure</a>  |



## Isotopic distribution cross-sections in inverse kinematics

| Beam | Target | Energy [A MeV] | Laboratory   | Reference   | Data  |
|------|--------|----------------|--------------|---|---|
| Fe   | H      | 300            | GSI, Germany | C. Villagrassa-Canton et al., Phys. Rev. C 75 (2007) 044603 -   | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">GSI</a>   |
| Fe   | H      | 1000           | GSI, Germany | C. Villagrassa-Canton et al., Phys. Rev. C 75 (2007) 044603 -<br>P. Napolitani et al., Phys. Rev. C 70 (2004) 054607 -  | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">GSI</a>   |
| Pb   | H      | 500            | GSI, Germany | L. Audouin et al., Nucl. Phys. A768 (2006) 1 -  | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">EXFOR</a> |
| Pb   | H      | 1000           | GSI, Germany | T. Enqvist et al., Nucl. Phys. A686 (2001) 481 -  | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">GSI</a>   |
| U    | H      | 1000           | GSI, Germany | J. Taieb et al., Nucl. Phys. A 724 (2003) 413 -<br>M. Bernas et al., Nucl. Phys. A765 (2006) 197 -<br>M. Bernas et al., Nucl. Phys. A 725 (2003) 213 -<br>M. V. Ricciardi et al., Phys. Rev. C 73 (2006) 014607 - | <a href="#">data</a>   <a href="#">figure</a>   <a href="#">GSI</a>   |



# Benchmark of Spallation Models

## List of participating models

- ↳ CEM0303 (A. Gudima)
- ↳ CEM0302 (S. Mashnik)
- ↳ PHITS-jam (N. Matsuda)
- ↳ PHITS-Bertini (N. Matsuda)
- ↳ PHITS-JQMD (N. Matsuda)
- ↳ Cascade04 (H. Kumawat)
- ↳ Isabel-SMM (Y.Yariv / A. Botvina / D. Mancusi)
- ↳ Isabel-Gemini (Y.Yariv / R. Charity / D. Mancusi)
- ↳ Isabel-ABLA07 (Y.Yariv / A. Kelic / V. Ricciardi / D. Mancusi)
- ↳ Geant4-Bertini (D. Wright)
- ↳ Geant4-BIC (D. Wright)
- ↳ Cascade-ASF (A. Konobeyev)
- ↳ CASCADEX (Y. Korovin)
- ↳ INCL4.5-SMM (J. Cugnon / A. Boudard / A. Botvina / D. Mancusi)
- ↳ INCL4.5-Gemini (J. Cugnon / A. Boudard / R. Charity / D. Mancusi)
- ↳ INCL4.5-ABLA07 (J. Cugnon / A. Boudard / A. Kelic / V. Ricciardi / D. Mancusi)
- ↳ FLUKA\* (A. Ferrari) \* Request to participate and send results in November

# Benchmark of Spallation Models

## Objectives

- To assess the prediction capabilities of the spallation models used or that could be used in the future in high-energy transport codes
- To understand the reason for the success or deficiency of the models in the different mass and energy regions or for the different exit channels
- To reach a consensus, if possible, on some of the physics ingredients that should be used in the models.




# Benchmark of Spallation Models

## Objectives

➤ To assess the prediction capabilities of the spallation models used or that could be used in the future in high-energy transport codes

➤ Direct visual comparisons between data and calculation

➤ Figures-of-Merit / deviation factors



# Benchmark of Spallation Models

## Objectives

- To assess the prediction capabilities of the spallation models used or that could be used in the future in high-energy transport codes
  - not to define “the best” model but for each observable, for different energy or mass range, give recommendations to use one model rather than another one
  - Report for end of November

# Benchmark of Spallation Models

## Objectives

- To understand the reason for the success or deficiency of the models in the different mass and energy regions or for the different exit channels
  - To reach a consensus, if possible, on some of the physics ingredients that should be used in the models.
- 
- Analysis of additional information, discussion with model authors
  - Second Meeting in February 2010
  - Presentation of the conclusions at ND2010

# Additional information on the calculation

- model  $\sigma_R$  ( $\sigma_{\text{geom}} \times N_{\text{inel}} / N_{\text{evts}}$ )
- $\sigma'$  used (normalisation)
- $E^*, E^*/A_R, A_R, Z_R, P_R, J_R$  distributions to enter de-excitation
- Multiplicities of n, p,  $\pi$ , lcp, IMFs from 1st stage and de-excitation

→ To understand the reasons for success or deficiency

# Benchmark of Spallation Models

## Possible continuation ?

- A “dynamical” continuous benchmark so that end-users of spallation models in transport codes have up-to-date information
  - new versions of the models / new models compared to the benchmark set of data added on the website
  - new experimental data : ask authors to do additional calculations or do calculations with the version of the code given by the authors

# Agenda Tuesday October 6th

|             |   |   |
|-------------|---|---|
| 9:00-9:15   | Welcome   | G. Mank (IAEA)  |
| 9:15-9:45   | Recall of benchmark objectives  | S. Leray (CEA – France)                                 |
| 9:45-10:15  | Technical aspects   | J.-C. David (CEA – France)                              |
| 10:15-10:35 | Break   |   |
| 10:35-11:05 | Result production and web site tools                                      | M. U. Khandaker (IAEA)                                  |
| 11:05-11:50 | Discussion on result production and web site                              | All participants<br>Convener: J.C. David (CEA- France)  |
| 11:50-12:10 | Expected outcome on the physics of spallation models                      | Y. Yariv (Soreq – Israel)                               |
| 12:10-12:30 | Processing of benchmark data  | J. Yoo (KAERI, Korea)                                   |
| 12:30-14:00 | Lunch   |   |
| 14:00-14:30 | Deviation factors   | A. Konobeev (FZK – Germany)                             |
| 14:30-15:00 | Recent progress in metrology and its statistical foundations              | R. Michel (ZSR – Germany)                               |
| 15:00-16:00 | Discussion on Deviation Factors / Figures of Merit                        | All participants<br>Convener: Y. Yariv (Soreq – Israel) |
| 16:00-16:30 | Break   |   |
| 16:30-16:50 | Some remarks about the experiments on neutron, LCP, and pion measurements | D. Filges (FZJ – Germany)                               |
| 16:50-18:00 | Results: General trends (Neutron – lcp – pion)                            | All participants<br>Convener: D. Filges (FZJ – Germany) |

# Agenda Wednesday October 7th

|             |  |   |
|-------------|--|---|
| 9:00-9:10   | Lessons from the former benchmark on spallation residues | R. Michel (ZSR – Germany)                               |
| 9:10-10:30  | Results: General trends (Residue – Excitation function)  | All participants<br>Convener: D. Filges (FZJ – Germany) |
| 10:30-11:00 | Break  |   |
| 11:00-11:30 | The view point of users                                  | F. Gallmeier (ORNL, USA)                                |
| 11:30-12:00 | Interaction with end-users                               | All participants<br>Convener: G. Mank (IAEA)            |
| 12:00-13:00 | Results: Model analysis - Strategy/Methodology           | All participants<br>Convener: G.Mank (IAEA)             |
| 13:00-14:00 | Lunch  |   |
| 14:00-15:00 | Conclusion & next steps                                  | All participants<br>Convener: S. Leray (CEA)            |
| 15:00-16:00 | Preparation of report and executive summary              | All participants<br>Convener: G.Mank (IAEA)             |